

Index

- Amorphous organic matter
 - aquatic origin, 280
 - identification, 275–87
 - land-plant origin, 275, 299
 - types, 275, 280
- Basin evolution
 - Aquitaine, 223
 - extensional, 253
 - Faeroe–Shetland, 223
 - heat flow, 253
 - and hydrocarbon generation, 232, 253–4
 - Lusitanian, 223
 - Molasse region, 232
 - N. Ireland, 216–18
- Basin—exploration success (UKCS), 357–71
- Basin modelling, 223, 232, 237–52, 253
- Beatrice Oil Field, 35–50
 - geological setting, 36–7, 42
 - source rocks, 147–59
 - surface prospecting exploration, 35–50
- Barents Sea
 - Barents shelf geological setting, 86
 - exploration, 90
 - structural elements, 91
- Biological markers
 - biodegradation, 191
 - Brae oils, 210
 - maturity, 155, 175–93, 242, 253
 - oil composition, 175–93, 210, 237–52, 253
 - porphyrins, 309–15
 - Upper Jurassic, Inner Moray Firth, 147–59
- Brae Fields, 203–12
- Carboniferous sediments
 - Barents Sea, 88
 - C. England, 67
 - E. Midlands, 67
 - history, 262
 - maturity, 28, 229
 - N. Ireland, 213–22
 - N. Sea Coals, 255
 - onshore UK, 67
 - stratigraphy, 217
- Carbon isotopes
 - gas, 227, 229
 - gases Central Europe, 229
 - geochemical prospecting, 51–63
 - kerogen, 209
 - of carbonate cements, 140
 - oils, 209, 211
 - sandstone reservoirs, 127–45
- Carbonate cement, location of, 143
- Celtic Sea, 223
- Central North Sea
 - burial diagenesis, 161–72
 - surface geochemical studies, 59–62
- Chemical weathering, 116–23, 274
 - redox and acid reactions, 117
- Clay minerals, abundance/variation, 163–4
- Cleveland basin, 29–33
- Coal
 - brown coal, 267–74
 - coalification, 267–74
 - extracts, 169
 - gas, 227, 229
 - measures, depth and maturity, 21, 27, 33, 229
 - N. Ireland, 216–18
 - oil potential, 255–65
 - porosity, 259
 - properties, 255–65, 267–74
 - pyrolysis, GC 263, 267–74
 - rank, 258
 - vitrinite reflectance, 296, 304
- Department of Energy, 3–9
 - divisions, 3
 - function, 3
 - geological and geophysical, 3–9
 - geological information, 319, 343–56
 - onshore, 67
 - organisation, 4
 - well record centre, 6, 356
- Diagenesis
 - burial, 163, 175
 - differentiation, 115
 - history of sandstones, 113–24, 127–45
 - kinetics, 115
 - mudstones, 121
 - reactions, 116, 122, 135, 139, 163
 - sequence, 127
- Drilling, UKCS exploration, 361–71
- Earth Science Bibliography
 - databases, 322, 327–8, 329–41
 - exploration success, 357–71
 - hydrocarbon shows, 341
 - North Sea, 321–8, 329–41, 357–71
 - production, 325
- Earth Science Literature
 - information gathering, 322, 329–41
 - UKCS, 321–8, 329–41, 357–71
 - UKCS exploration success, 357–71
- East Shetland Platform, 95–111
- Economic Commission for Europe (ECE),
 - Committee on Gas, 11–17
- English Channel, 223
- Europe
 - gas exploration, 11–17, 19–33
 - gas reserves, 11–17
- Exploration
 - and basin modelling, 237–52
 - and geological information, 317–73
 - bibliography, 321–8
 - data acquisition and release, 6, 343–56
 - gas, 12–17, 19–33
 - geochemical, 35
 - N. Ireland, 213–22
 - North Sea, 69, 343–56
 - North Sea Plays, 73–82, 174–94, 195–201, 202
 - Norwegian Shelf, 71–93
 - offshore, 17

- onshore UK, 67–8
- Po Valley, Italy, 227
- porphyry studies, 309–15
- released wells, 346
- surface prospecting, 35–50, 51–63
- UKCS history, 361–71
- UKCS information, 343–56
- UKCS success, 357–71
- Western and Northwest Europe, 223–4
- Exploration boreholes
 - German Alps, 231
 - N. Ireland, 218
- Facies distribution
 - Moray Firth, 195, 200
 - North Sea, Lower Jurassic, 80
 - organic facies, 275–87
 - source rocks, 190, 195, 200
 - South Viking Graben, 195, 200
 - Upper Jurassic, 195–200
- Field development plans
 - Europe, 12
 - UKCS, 12
- Fladen ground spur, 95–111
- Fluorescent studies
 - spectral correlation, 289–94
 - surface prospecting, 38–50
- Forties field, surface prospecting exploration, 60–2
- Gas
 - accumulations, 24, 229, 361, 363–6
 - biogenic, 227, 229
 - coals, 229, 255–65
 - composition, 22, 25
 - deep horizons, 13, 229
 - discovery in N. Ireland, 213
 - exploration, 11–17, 19–33, 227, 363–6
 - generation, Po Valley, 227
 - occurrence, Po Valley, 227
 - reserves, 11–17, 19, 69, 73, 227, 363–6
 - reservoirs, 19–33, 227
 - source rocks, 19–33, 227, 229
- Gas chromatography, 151, 169, 203–12, 239–46, 258–65, 267–74, 299
- mass spectrometry (GCMS), 155, 210, 239, 267–74
- Generation
 - basin evolution, 253–4
 - basin modelling, 237–52, 253
 - Central Europe gas, 229
 - from coals, 255–65
 - hydrocarbons, 147–59, 175–93, 195–200, 203, 255, 295, 309
 - material balances, 248
 - peak, 78, 181, 259
 - Po Valley gas, 227
 - porphyrins, 309
 - timing of, 29, 78–83, 175–93, 200, 314
 - vitrinite reflectance, 295
- Geochemical log, 244
- Geochemical studies
 - basin modelling, 237–52, 253
 - coals, 255–65
 - German Alps, 231–3
 - Inner Moray Firth, 147–59
 - isotope studies, 227, 229
 - Norwegian source rocks, 73–93
 - NW Europe Gas Province, 19–33
 - parameters, 176, 253, 309
 - porosity enhancement and reduction, 113–24
 - porphyrins, 309–15
 - sandstone reservoirs, 127–45
 - screening methods, 237
 - shallow sediments, 51, 65
 - South Viking Graben, 161–72, 195–200, 203–12
 - surface prospecting, 35–50, 51–63, 65
 - techniques, 163, 175, 198, 237–52, 309
- Geological Databanks (UKCS), 329–41
- ERICO, 330
- petroleum information, 330
- UKCS information, 355
- UKCS success, 357–71
- use of computerized information, 373
- Geological Information, 317–73
- databanks, 329–41, 357–71
- exploration (UKCS), 343–56, 357–71
- files, 332
- hydrocarbon shows, 341
- UKCS well catalogue, 353
- use of computerized information, 373
- Geological Survey of Northern Ireland, 219
- Geothermal Studies, 219–21
- German Alps, 231–3
- Molasse region, 231
- Haltenbanken (Norway), 71–93
- exploration, 85
- High Pressure Liquid Chromatography, 309–15
- Hydrocarbon Fields, 343–56
- Beatrice Field, 37, 147
- Brent Field, 361
- Forties Field, 51, 62, 95, 361
- Frigg Field, 361
- Hewett Field, 33
- Montrose Field, 95, 361
- Statfjord Field, 72
- West Sole Field, 361
- Inner Moray Firth, 35–50, 147–59
- exploration, 361
- geology and structure, 149
- Upper Jurassic sediments, 147–59
- Institute of Geological Sciences (IGS), 5–9, 355
- nomenclature, 98, 99
- Jurassic Plays
 - Haltenbanken prospects, 86
 - Inner Moray Firth, 147–59, 198
 - Lower Jurassic play types, 88
 - Middle Jurassic play types, 83
 - porosity/depth trends, 85
 - South Viking Graben, 161–72, 195, 198
 - Upper Jurassic play types, 84, 90, 147–59
- Jurassic Sand Play
 - reservoirs, 127–45
 - Statfjord-Troll, 80
 - South Viking Graben/Central Graben, 161–72, 195

- Kerogen**
 amorphous, 275–87
 basin modelling, 237–52
 composition, 197, 199, 203, 255–65, 267, 287, 289
 facies, 147, 167, 195–200, 255, 275–87
 maturity, 150, 167, 175–93, 203–12, 237, 267, 289
 methods, 148, 163, 267, 275, 289–94
 occurrence, 77, 195–200, 203, 275
 oil potential, 78, 148, 175–93, 195–200, 255, 275, 314
 spectral correlation of SCI, 289–94
 type, 77, 150, 168, 175–93, 195–200, 203–12, 232, 287
- Kimmeridge Clay Formation**
 facies, 190
 Inner Moray Firth, 147–59, 198
 maturation, 173–93, 203–12
 oil discovery, 68, 148, 175, 195, 203
 source rocks, 76, 148, 159, 175–93, 203–12, 258
 South Viking Graben, 161–72, 198, 203–12
- Licencing**
 N. Ireland, 213
 Norwegian, 71
 On-shore, 67–8
 U.K., 5, 343, 353
- Light hydrocarbons**
 analysis, 229, 239–48, 258
 coals, 259
 isotopes, 229
 molecular composition, 51, 65, 239–48, 259
 surface prospecting, 35–50, 51–63, 65
 vitrinite, 299
- Lower Triassic Plays**
 deltas, 255
 play types, 89
- Lower Tertiary Sand Plays, Frigg, 78, 81**
- Macerals, 261, 267–74, 289–94, 295**
 analysis, 303
 identification, 275, 298
- Marine sediments, environment, 136**
- Maturation**
 assessment, 153, 180–93, 195–200, 237–41, 289, 295, 306
 amorphous organic matter, 275–87
 basin modelling, 237–52
 biomarkers, 155–9, 175–93, 309–15
 carboniferous, 29–32, 216
 Central Graben, 79
 Cleveland basin, 31
 coals, 257–65
 controls, 181, 199, 237, 248
 German Alps, 231–33
 Inner Moray Firth, 36, 150, 195
 Jurassic, 77, 78, 199
 organic matter, 118, 150, 167, 195, 227, 237, 248, 275
 porphyrins, 309–15
 scales, 289–94, 295
 source rocks, 51, 167, 175–93, 195–200, 227, 237–52, 275, 295
 time and maturity, 186, 200, 249–50, 295
 vitrinite reflectance, 295–307
- Mehakam Delta, coals, 255–65**
- Mesozoic basins**
 Ireland, 215
 Molasse region, 232
 Norway, 73–93
 S. England, 67
- Methylphenanthrene Index (MPI), 243**
- Microbial reactions**
 degradation, 118, 175, 195
 fermentation, 139
 oil, 191, 195
 organic matter, 276, 299
- Midland Valley of Scotland, 21, 68, 215–16**
- Migration, 187, 200, 231, 246**
 from coals, 255–65
- Mineral-porewater reactions, 114–15, 123, 142**
- Mudstones**
 diagenesis, 121
 evolution, 120
 sandstone links, 123
- National Environment Research Council (NERC), 5**
- Northern Ireland**
 geological background, 213
 geological map, 215
 gravity and aeromagnetic, 218
 search for petroleum, 213–22
 seismic survey, 220
 wells drilled, 214
- North Sea Plays, types, 73–93**
- Norwegian Petroleum Directorate, 71–93**
- Norwegian Shelf**
 exploration, 71–93
 Haltenbanken, 71–93
 Licencing, 71
 Traeabanken, 71–93
 Tromsi, 71–93
- Oil**
 biodegradation, 191
 composition, 175–93, 195, 197, 198, 200, 205, 245
 expulsion from coals, 261
 from coal, 255–65
 generation, 77, 177, 181, 195–200, 237, 261, 276, 295
 German Alps, 232
 kerogen type, 195–200, 237, 275
 migration, 187, 200, 231, 261
 North Sea reserves, 69, 357, 367–71
 Norwegian reserves, 73, 78, 357
 properties, 175–7, 198, 205
- Onshore U.K. exploration, 67–68, 343–56**
 location of well, 348
 released wells, 348
- Outer Moray Firth Basin, 95–111**
 hydrocarbon occurrences, 196
 kerogen types, 195–200
 oil generation, 195–200
- Oxygen isotopes, sandstones/carbonates, 127–45**
- Palaeocene Sands**
 depositional history, 105–109
 isopach map, 97

- lithostratigraphic scheme, 98–105
 - South Viking Graben, 161–72
 - stratigraphy and sedimentation, 95–111
- Paris Basin, 253, 309
- Permian sediments
 - N.E. England (reservoirs), 67
 - S. Gas Province, 22, 361
- Porphyryns, 309–15
 - maturation indicators, 309–15
 - oils, 311
 - sediments, 312
- Pyrolysis studies, 237–52, 255–65, 267–74, 314
 - coals, 255–65
 - mass spectrometry, 267–74
 - Rock-Eval, 237, 252, 255, 265, 314
 - vitrinite, 299
- Quantitative Basin Analysis, 246–52
 - conceptual model, 248–9
 - flow of information, 248
 - mathematical model, 246
 - maturity and hydrocarbon generation, 247
 - procedures, 248
- Resin (fossil), 272
- Sandstones
 - carbonate cements, 144
 - diagenesis, 122
 - diagenetic history, 113–24
 - Lusitanian Basin, 223
 - mudstone links, 123
 - N. Ireland, 221
 - primary and secondary porosity, 113–24, 127
 - reservoirs, 95–111, 127–45, 232
- Sedimentation
 - environments, 118
 - North Sea Palaeocene Sands, 95–111
 - Sandstone reservoirs, 127–45
- Sedimentary organic matter
 - amount, 76, 148, 150, 168, 203
 - coals, 255–65, 267–74
 - components, 119, 150, 168, 195, 201, 239, 267
 - identification, 275–87
 - Inner Moray Firth, 147–59
 - maturity, 150, 168, 175–93, 195, 203, 237, 253, 267
 - Norwegian Source Rocks, 71–93
 - porphyryns, 309–15
 - preservation, 286
 - reactions, 139
 - type, 77, 148, 195, 201, 239
 - Viking Graben, 166–9, 175, 195, 203
- Source rocks
 - amorphous organic matter, 275–87
 - basin studies, 186, 232, 237–52
 - Beatrice Oil Field, 36, 147–59
 - Brae oils, 203
 - Carboniferous, 19–33
 - coals, 255–65
 - and exploration, 62, 275–87
 - gas, 19–33
 - gas accumulations, 22
 - German Alps, 231–33
 - Inner Moray Firth, 147–59
 - maturity, 28–9, 147–59, 175–93, 236, 309
 - N. Ireland, 213, 221
 - Norwegian Region, 76–93
 - oil composition, 175–93, 309–15
 - oil correlation, 147–60, 161–73, 174–94, 195–201, 201–212, 232, 239–52, 309–15
 - Po Valley, Italy, 227
 - Viking Graben, 175, 195, 203
- Steranes, 155–9, 173–95, 210–11, 243, 253–4, 314
 - activation energy, 253
- Stratigraphy
 - Beatrice Field, 42
 - Cleveland Basin, 31
 - Haltenbanken, 87
 - N.W. Europe Gas Province, 19–22, 24
 - N. Ireland, 213–18
 - North Sea Palaeocene Sands, 95–111, 98
 - Troms I, 92
 - stratigraphic correlation (Palaeocene), 99
 - South Viking Graben, 163–4, 166
 - UKCS released wells, 350
- Structural features
 - basin evolution, 253
 - Barent Sea Shelf, 86, 91
 - Beatrice Field, 42
 - Brae area, 204
 - German Alps, 231–3
 - Inner Moray Firth, 147, 149
 - Mid-Norwegian Shelf, 82–5
 - More-Lofoten Shelf, 85
 - N.W. Europe Gas Province, 22, 31
 - N. Ireland, 213–22
 - N. Ireland and Scotland, 216
 - North Sea, 63
 - Palaeocene sediments (UK), 95–8
 - South Viking Graben, 161–2, 196, 204
- Surface geochemical exploration, 35–50, 51–63, 65
 - laboratory studies, 52, 55
 - method, 36, 52, 55, 65
 - pattern recognition, 38, 53, 56
 - 'Sniffer' method, 65
 - statistical methods, 39, 60
- Traenabanken (Norway), 71–93
 - exploration, 85
- Transmitted light microscopy
 - amorphous organic matter, 275–87
 - equipment, 291
 - spectral correlation of SCI, 189–94
 - spore colour, 195–201, 289–94
 - TAI, 237–52, 289–94
- Triterpanes, 155–9, 173–95, 210–12, 253–4, 314
 - activation energy, 253
- Troms I (Norway)
 - prospects, 92
 - stratigraphy, 92
 - well results, 92–3
- Tromsøfket (Norway), 71–93
 - exploration, 85, 90
- Upper Cretaceous Plays, Danian play types, 82, 92
- Viking Basin (North Sea), 96, 110

- Viking Graben (North Sea)
 - clay reactions, 164
 - exploration, 361
 - maturation, 170, 195–200
 - oil generation, 159, 161–72, 195–200
 - tectonic setting, 95–111, 161, 196
- Vitrinite reflectance
 - basin studies, 239–52, 253, 314
 - Cleveland Basin, 30
 - coals, 256–65, 267–74
 - German Alps, 231
 - Inner Moray Firth, 148, 151–5, 197
 - maturity tool, 231, 295–307, 314
 - measurements, 302
 - methods, 295–307
 - sample preparation, 295
 - South Viking Graben, 166, 169, 197
 - Southern North Sea, 33
 - types, 295–307
 - vitritine, 267–74, 276
- Witch Ground Graben
 - exploration, 361
 - tectonic setting, 95–111
- Wood, 272
- UKCS Exploration History, 361–71