

# Index

References in *italics* are to Figures or Tables

- Addington, Croydon, tracer tests to confirm pollution source 98
- adits, built into the Chalk aquifer 99–100
- aggressive water, and karst features 104
- agriculture, in wastewater irrigation areas 156
- ammonium 55
- anions, major 55
- apparent ground conductivity measurements 80–1, 90
- apparent conductivity maps 83, 85, 86, 88
- interpreted resistivity models 82
- low tidal cycle 82, 84
- seawater incursion and saline intrusion 83, 86
- time/tide dependent experiment 81
- aquifer evaluation methods, Ogallala Formation 64–5, 66, 70
- aquifer pollution, anthropogenic 61
- aquifer pollution vulnerability 7
- mapping of 9–12
- systems in use in Britain 10–12
- aquifer vulnerability maps 2, 3, 19–20
- aquifers 211
- British, recharge and pollution vulnerability 7–22
- Chalk, London Basin 47–62
- classified, England and Wales 192–3
- dual-porosity 2
- vulnerability of 12–16
- High Plains Aquifer 63–70
- Irish, classification of 181
- Jersey, nitrate contamination 23–35
- karstic, problems in calculating GPZs 202–3
- perched 45, 157
- Sherwood Sandstone Group 191, 192, 202
- susceptibility to side-effects of groundwater exploitation 71–6
- Tawilah Sandstone 169
- unconfined 202
- nitrate in 3
- vulnerability and recharge 3–4
- ‘aquiperm’ 11
- aquitards 11, 130
- leakage from 72, 75
- areal water balance method 111
- Arish Mell, Dorset, tracer experiment 98
- attenuation
- by matrix component, vadose zone flow 16
- by matrix diffusion 18
- of contamination, Ireland 181, 183
- of pollutants, in the vadose zone 8–9
- processes to prevent pollutants reaching groundwater 168
- autotrophic denitrification 31, 34
- bacteria, in groundwater 13, 102, 102, 103–4
- Barton Beds 77
- Basal Sands
- dewatering of 49
- hydraulic continuity with the Chalk 48
- leaching from 61
- sulphide oxidation when dewatered 56
- Beachy Head cave system 98–9
- bedrock topography, and saline intrusion 86, 89
- best-estimate zone (BEZ) 208
- Blisworth Clay 214
- Blisworth Limestone 214, 216
- boreholes
- in the Chalk 99, 100, 100
- Chilgrove borehole 100, 118
- Jersey 143
- Léon, Mexico 157, 165
- Redgrave borehole, reducing discharge of calcareous water 129
- boron 55, 57, 61
- boulder clay, cover over Waveney and Little Ouse catchments 139
- Bridget’s Farm, Hants
- fracture flow 119
- matrix conductivity 127
- Brighton water supply, adit system for 100
- Brule Clay 64
- by-pass flow 117, 118
- Bridget’s Farm and Fleam dyke 119
- controlling variables 121–2
- controls on onset 127
- importance increases with rise of matrix potentials 118–19
- linked to significant recharge events 123
- and soil moisture deficit 122–3
- by-pass routes, role of 3
- calcite dissolution 49, 51, 61
- canals, unlined 112
- carbonate chemistry, London Basin groundwater 55
- carbonates, incongruent solution of 59, 60, 61
- Carboniferous Limestone 193
- groundwater protection zones 207
- catchments
- hydrology, Redgrave area 133–6
- mixed, Jersey 148, 149, 150, 151
- cation exchange 61, 195
- cations, major and minor 55, 56
- caves/cave systems, in Chalk 98–9
- cesspits, for sewage disposal 174
- Chalk 47, 191, 192
- as bedrock beneath coastal wetlands 79–80, 82, 83, 89–92
- contrasting catchments, near Redgrave Fen 129–41
- fracture apertures and spacings 127
- hydraulic properties 94
- hydrochemistry
- evolution of, London Basin 49, 50, 51, 51
- influence of pore water on 57, 59
- karstification in 4
- porosity sets 2
- preferential vadose zone flow 14–15, 19
- rapid groundwater flow in 95–106
- salt concentrations in 83
- surface drainage 100
- transmissivity, London Basin 49
- Chichester flood, role of groundwater 100
- chloride 57, 61
- chloride balance, confirms recharge from wastewater 159

- chloride concentrations  
 Kellaways Sand 215  
 Léon, Mexico 162, 163, 166, 167
- chloride enrichment 51
- chromium 155  
 not reaching the water-table 164
- Clay-with-Flints 16, 19
- clays, Jurassic 193
- climate change, effect of 20
- coastal sediments, affected by in-phase tidal movement 89, 92
- coastal wetlands  
 environment 77–8  
 geophysical investigation, saline intrusion/geological structure 77–94  
 data processing and presentation 82  
 empirical models 89–92  
 environmental implications 92  
 geophysical field methods 80–1  
 presence of channel features 83, 86, 89
- cones of depression, Léon, Mexico 160, 162
- contaminant breakdown, vadose zone 16
- contaminants, Ireland  
 quantity of 183  
 release points of 182–3
- contamination  
 concept of vulnerability to 182  
 intermittent, with faecal bacteria 13  
 risk to groundwater 180  
 source control, groundwater protection 188
- continuous coring, for an accurate geological model 40
- Crag 193
- Cuatralba Ignimbrite 156, 157
- deglaciation, Devensian 44–5
- denitrification 16, 18  
 Jersey 24, 31–3, 34
- dewatering, and effective stress 72
- diagenesis, induced in aquifers 73
- direct recharge 107, 108, 109
- dispersive flow *see* preferential vadose zone flow
- dolines 101–2, 103
- DRASTIC methodology (US) 12, 181, 191
- drift geology 39  
*see also* boulder clay
- Drigg low-level waste facility 37–46  
 location and description 37–9
- drilling, rotary and cable percussion, Drigg site 40–1
- dry valleys, Chalk 100, 101
- East Anglia, non-karstic tracer tests 98
- EC, Groundwater Directive 211
- effective rainfall-river transfer function  
 modelling 135–6, 136, 137, 140
- effective stress 72
- engineered containment  
 costs of 211–12  
 not necessary in Lower Oxford Clay 216
- Environment Agency, delineating groundwater protection zones 201–6
- ephemeral channels, recharge from 112
- Escherichia coli* *see* bacteria
- exploitation 47  
 chronology of impacts 75  
 indicators of side-effects 72–3  
*see also* over-exploitation
- faults 23
- Fe concentration, at reducing groundwater sites 32
- fissures, large, preferential flow in 13
- Fleam Dyke lysimeter case study 117–28
- FLOWPATH GPZ program 204, 205, 208
- forest precipitation interception experiment, Drigg 42
- fracture flow *see* by-pass flow
- fractures 3, 10, 95, 117–18  
 aperture related to matric potential 126  
 dilated 23  
 dissolution-enhanced 104, 105  
 ease transport of *E. coli* 102  
 rapid infiltration through 118
- Gault Clay 48
- Geological Survey of Ireland, Quaternary geology maps 185
- geophysical surveys  
 Drigg 41  
 saline intrusions, coastal wetlands 80–2, 92
- glacial tills 2, 16  
 Irish, need to determine permeabilities 187  
 over-compacted, Drigg site 43–4
- GOD systems, for vulnerability index estimation 10, 11
- GPZs *see* groundwater (source) protection zones
- Great Oolite Group 214
- groundwater abstraction  
 Léon, Mexico 159  
 exceeding recharge 160, 167  
 London Basin, altering natural flow directions 48–9, 61
- groundwater chemistry  
 changes in, London Basin 47–62  
 influence on Chalk hydrochemistry 65–6, 70
- groundwater movement, tide-controlled, shallow saline and fresh 90, 90
- groundwater pollution *see* pollution
- groundwater protection 4
- Groundwater Protection Policy (GPP) 191  
 disposal of waste materials 211  
 key elements and variables 191–2  
 Major, Minor and Non-aquifers 211  
 maps, vulnerability system 191
- groundwater quality  
 deterioration in 61, 73, 74  
 Léon, Mexico 160–5  
 not addressed by vulnerability maps 197  
 poor, Kellaways Sand 215, 216
- groundwater resource development, Nebraska 63–70
- groundwater source 199  
 relationship with protection zone 200, 200
- groundwater (source) protection zones 199–210  
 definition of 199–200  
 delineation methods  
 guidelines needed for use of other models 209  
 history of 200–1  
 improvement needed in some areas 208–9  
 selection of 206–7  
 used by Environment Agency 201–6

- matrix of 188, 188, 189
- model of flow system surrounding source 206–7
- precision and uncertainty in delineation 207–8
- groundwater store 147, 148
- groundwater vulnerability *see* vulnerability
- groundwater zoning
  - and diffuse-source pollution 197
  - London Basin 49, 51, 59, 60, 61
- groundwater(s)
  - calcareous, reduced discharge of 129
  - decline in level of 72, 74, 75
  - is protection worthwhile? 211–17
  - Jersey, nitrate pollution of 23–35
  - and nitrate concentrations 3
  - perched 40
  - possible impacts of wastewater use 153
  - reducing, increasing concentrations of SO<sub>4</sub><sup>2-</sup>s 32
  - saline, interpreted channel-limited migration of 90, 91, 92
- Hampshire syncline 79
- Havant-Bedhampton tracer experiment 97
- hazard 180
- heterotrophic denitrification 31, 32
- hydraulic conductivity 173
  - in Lowestoft Till 17
  - of the matrix, Fleam Dyke 122
  - rapid rise in 119
- hydraulically effective rainfall (HER) 130, 135–6, 137
- hydrochemical facies, London Basin 56
- hydrochemistry
  - changes induced due to water-rock reaction 73
  - London Basin, regional trends 49–51
- hydrological model, Trinity catchment
  - description 146–8
  - instrumentation for data 144–5
  - results 148–9
  - strategy 146
- hydrologically effective precipitation (HEP) 125–6, 128
- hydrostratigraphy, Drigg, interpreted 42–5
- impermeability, of Kellaways Sand 215
- incongruent solution 59, 60, 61
- indirect recharge 107, 109
- infiltration
  - from groundwater irrigation 160
  - from wastewater irrigation 170
  - of surface flows 170–2
  - travel time of 183
- infiltration flux 121
- integrated vulnerability mapping 10
- intensive farming, and nitrate pollution, Jersey 27, 34
- intergranular flow 94
- intergranular porosity 23
- intergranular seepage 118
- ion exchange 51, 57, 58, 61
  - reverse ion exchange 61
- ions, major and minor, analysis of 52–3
- irrigation
  - Léon, Mexico
  - from groundwater 159–60
  - wastewater 153–6, 159, 168
  - water from upper part of aquifer 167
  - rice, and recharge 112
- Sana'a Basin, Yemen 169
- recharge to groundwater from 172–3
- water for, Jersey 143
- Jersey
  - estimation of groundwater recharge 143–52
  - nitrate pollution of groundwaters 23–35
  - abstraction concentrated in upper bedrock aquifer 33, 34
  - deeper groundwater component 33, 34
  - hydrochemical analyses from bedrock aquifer 24–8, 29–30
- karst
  - in Chalk 96–7
  - karst-like solution features 15–16
- karst areas, vulnerability mapping 185
- karst-type behaviour, Chalk of southern England 96–102, 103–5
- karstic limestone 2, 3–4
  - see also* Carboniferous Limestone
- Keele Beds 202, 203
- Kellaways Clay 214
- Kellaways Formation 212
- Kellaways Sand 4
  - corresponds to Non-aquifer definition 216
  - geology of 212–14
  - hydraulic conductivity data 214–15, 214
  - hydrogeology 214–15
  - petrographic analysis of 214
  - seepage rates to R. Ouse 216
- Kilham, N Yorks, tracer test 98
- land-surface zoning 9, 188
- landfill
  - engineered containment 211
  - in the Lower Oxford Clay 212, 213
- Langstone Harbour, UK, geophysical investigation of coastal wetland 77–94
- Lea Valley, rainfall-bacteria level correlation 97
- leaching potential classification 195–6
- leather industry, polluting effluent 154–5
- Léon, Mexico
  - evolution of groundwater system 160, 162
  - groundwater quality 160–5
  - groundwater resources 158–60
  - natural sources of recharge 157
  - options for maintaining aquifer quality 166–7, 166
  - security of potable supplies 166–7
  - wastewater irrigation system 153–6, 159
  - concern over pollutants in 166
- liner costs 211–12
- Little Ouse River 131–3
  - boulder clay catchment cover 139
  - river flow hydrograph analysis 134–6, 137
- London Basin
  - equilibration of pore-water with fissure water 57, 59
  - exploitation of Chalk aquifer 47
  - geology 47–8
  - groundwater zonation 49, 51, 59, 60, 61
  - hydrogeology 48–9
- London Clay 48
  - ion-rich leakage from 56
  - leaks calcium and magnesium sulphates 51

- low-level waste (LLW) site characterization,  
 Drigg 37–46  
 existing radiological safety case 39  
 geological and hydrogeological investigation  
 techniques 40–2  
 hydrostratigraphy 42–5  
 water balance 42, 43
- Lower London Tertiaries (LLT) 48  
 Lower Oxford Clay 212, 216  
 Lowestoft Till, protection of underlying aquifers 16–18
- M1/M25 intersection, tracer experiment 98  
 Magnesian Limestone aquifers 191, 192  
 magnesium 51, 59, 60, 61  
 matric hydraulic conductivity 118  
 matric potential 118, 121, 126  
 high, and drainage peaks 122  
 relationship with hydraulic conductivity 119  
 matrix flow 117, 122  
 Mercia Mudstone Group 193  
 microfissures and macrofissures 118  
 microfractures, dominating flow 98  
 Middle Chalk, water transport through 117–28  
 MODFLOW and MODPATH (USA), used in GPZ  
 programs 204, 206  
 Mole, River, swallow holes 102, 104  
 mountain front recharge 112
- $^{15}\text{N}$  isotope tracing technique 24–8, 29–30  
 $^{15}\text{N}$  trends  
 denitrification 31–3  
 mixing between end-member groundwaters 33  
 spatial variation in source of dissolved  $\text{NO}_3^-$  28, 31  
 nitrate, in unconfined aquifers 3  
 Nitrate Sensitive Areas: Nitrate Vulnerable Zones 208  
 non-aqueous phase liquids, dense (DNAPL) and light  
 (LNAPL) 7, 197
- Ogallala City, area physiography and geology 63–4  
 Ogallala Formation 63, 64  
 aquifer evaluation methods 64–5, 66  
 groundwater chemistry 65–6, 70  
 shallow and deep water zones 64–5, 66, 66, 67, 68, 69,  
 70  
 optimum vulnerability mapping 184–5  
 organic matter 16, 195  
 oxidation of 32, 34  
 Ormskirk Sandstone 39, 42  
 over-exploitation 4  
 concept of 71–2  
 over-irrigation 114, 173  
 Oxford Clay 213
- Palaeogene cover, and fast flow routes in Chalk 97–8,  
 103–4, 105  
 permanent water bodies, recharge from 112–14  
 permeability 193  
 secondary 23  
 pesticides, transport to aquifers 13, 14, 15  
 piezometric levels, Kellaways Sand 215  
 piston flow 118  
*Policy and Practice for the Protection of Groundwater*,  
 NRA 1, 199
- pollutants 3  
 assessment of migration within a wetland  
 environment 92  
 build-up of, Léon, Mexico 158  
 transport by preferential flow 13  
 in wastewater, Léon, Mexico 162, 164, 165
- pollution  
 diffuse-source 3, 185, 197  
 induced 73  
 management of potential sources of 3  
 nitrate 3  
 Jersey groundwaters 23–35  
 point-source 3, 197–8
- pollution risk  
 deep groundwaters, Léon, Mexico 161  
 defined 9–10
- pore-water  
 diffusion from 61  
 influence on Chalk hydrochemistry 57, 59  
 pore-water pressure, decreased by abstraction 72  
 Portsdown Monocline 79  
 potassium 61  
 precipitation, recharge from 109–11  
 preferential vadose zone flow 13–16  
 pumping rate/protected yield, in GPZ delineation 206  
 pumping test data, limitations of 95–6
- Quaternary deposits 197  
 variability of 193–4  
*see also* boulder clay; glacial tills; superficial deposits  
 quickflow store 147, 148
- radionuclides, transport within aquifers 40  
 rainfall-recharge-run-off model, Jersey 146–8, 147  
 rainfall-run-off model, Sana'a Basin 171  
 rapid groundwater flow 95, 96–102, 103–4  
 recharge 1–5, 107–15  
 and aquifer vulnerability 3–4  
 components of, Sana'a Basin 170–6  
 from intermittent flow 112  
 from permanent water bodies 112–14  
 from precipitation 109–11  
 in karstic areas 184  
 local 111  
 and pollutant transport, Léon, Mexico 153–68  
 and pollution vulnerability 7–22  
 through the drift, near Redgrave Fen 129–41  
 to Chalk aquifer, London Basin 48  
 to regional aquifer, Drigg LWW site 45  
 recharge distribution, modelling catchment  
 contrasts 137–40  
 recharge estimation 107, 108–9, 111, 114  
 challenges in 109–14  
 Jersey 143–52  
 lysimetric approach to 120  
 overall, Redgrave area 136–7  
 the pitfalls 1–2  
 in semi-arid areas 169–77  
 soil physics approach to 120–1  
 recharge evaluation techniques 7, 8  
 recharge events 118  
 timing/duration, Fleam Dyke 125, 127–8  
 recharge mechanisms 3, 7, 39, 117  
 recharge processes 107–8

- recharge spatial variability, assessment and prediction  
of 109, 111
- recrystallization, of Chalk 51
- Red Crag 129
- redox change, in aquifers 73
- redox reactions 61
- resistivity tomography surveys 81, 82, 83, 85, 88, 90
- risk 188  
and engineered containment systems 216–17
- risk management 180, 188
- river terrace deposits 193
- run-off 1–2  
rapid-response 148
- saline intrusion 73  
coastal wetland 79, 92  
distribution and geometry of 82–9  
in- and out-of-phase with tidal cycle 89, 92  
geological control on 89  
geophysical properties of 83, 89  
Jersey 23  
salt concentrations 89  
susceptibility of aquifers to 74  
tidal control on 89
- saline water, deep, upconing of 73
- salinity, of water from Kellaways Sand 215
- saltmarsh 77  
tidal control of sedimentation 77–8
- Sana'a Basin, Yemen, recharge estimates 169–77  
total recharge 176–7  
wadi recharge estimation 171–2, 172
- sediments, compacted by abstraction 72
- seepage recharge 112
- (semi)-arid areas  
recharge estimates 169–77  
recharge of phreatic aquifers 107–15
- septic tanks  
pollution by 3, 31  
recharge from 113, 159
- sewage  
leaking sewers, recharge from 113–14  
polluting groundwaters 28, 31  
used in wastewater irrigation 154, 159
- Sherwood Sandstone Group aquifers 191, 192  
Wilmslow Sandstone unit 202
- sodium 51, 57, 61
- soil 194–5
- soil hydraulic properties, indirect estimation  
methods 109
- soil maps, England and Wales 195–6
- soil moisture deficit (SMD), and by-pass flow 122–3,  
127
- soil moisture store 147–8
- soil water balance model  
Drigg 42, 44  
Redgrave area 133
- soil zone 12  
in the infiltration process 2, 9  
potential role of 194
- solution activity  
associated with cover 104  
*see also* incongruent solution
- solution features, karst-like 15–16
- solution pipes 100, 101–2  
associated with cover 103
- South Mimms, swallow holes 97
- stable depth 72
- stable isotopes, London Basin groundwater 53, 55, 56
- Stradbroke Depression 129
- strontium 51, 55, 59, 60
- sub-soils 183  
mapping of 190
- subglacial tunnel valley, drift-filled 129
- subsidence 72–3, 74
- sulphates  
London groundwater 56, 61  
Ogallala aquifer 66
- superficial deposits  
clay, effect of 16–19  
permeable/low-permeability deposits 194  
unconsolidated, as minor aquifers 193
- superficial deposits, mapping of 2
- surface leakage, enhanced 56, 61
- swallow holes 97, 101–2, 103, 104
- Tawilah Sandstone 169, 173, 175, 177  
groundwater levels 175
- Texas and New Mexico High Plains, recharge through  
playa basin floors 111
- tidal control, on saline intrusion 89
- toxic and trace metals, accumulation of 158
- tracer tests 103  
karstic and non-karstic 97–8
- transpiration, estimation of 1
- Trinity Valley catchment, Jersey  
model strategy, description and results 146–9  
monitoring network 143–6
- unsaturated flow, modelling of 173, 174
- unsaturated zone 184, 192
- Upper Bracklesham Group 77
- Upper Chalk, tritium movement through 118
- Upper Helsby Sandstone 202, 203
- urban development, impacts on groundwater  
recharge 109  
*see also* septic tanks; sewage
- urban recharge 112–13  
Sana'a Basin 173–6
- US Soil Conservation Service curve number  
method 170
- vadose zone 7–9  
contaminant breakdown 16  
environmental tritium in 13, 13, 14  
need to improve zone sampling and monitoring 20  
water movement in 8–9
- valleys, as sites of rapid groundwater flow 104
- vulnerability  
component of risk and risk assessment 180  
integration into the groundwater protection  
scheme 188–9  
'laws of' 10  
need for proper evaluation of risk 216–17  
site assessments of 187

- vulnerability assessments 180
  - England and Wales 191–8
  - Ireland
    - factors determining process 182–3
    - geological and hydrogeological parameters used 183–4
    - limitations and uncertainty in 186–7
- vulnerability concept 9–10, 179
  - criticisms of application of 189
  - criticisms answered 189–90
  - for risk assessment 187
- vulnerability mapping 9
  - England and Wales 191–2, 193, 195–6
  - Ireland 3, 179–90
  - guidelines for 184–6
  - procedural issues in 19–20
- vulnerability maps
  - England and Wales, and the non-specialists 197–8
  - Ireland
    - disclaimer on 186–7
    - not an alternative to site-specific investigation 190
    - scales 186
- vulnerability ratings, Ireland 185–6, 187, 188
- wadi flows 170, 171
- waste regulators, use of NRA aquifer classifications 211
- wastewater
  - from cesspits, in urban recharge 174–5
  - as a resource 153
- water balance
  - Drigg low-level waste facility 42, 43
  - Léon, Mexico **158**, 159
  - Redgrave and Lopham fens 133–4
- water balance model, Trinity catchment 146–9, 149–50
- Water End, Herts, swallow-hole activity 97, 104
- Water End-Lea Valley, rapid natural flows 97
- water mains, leakage from 113, 173
- Waveney, River 131–3
  - boulder clay catchment cover 139
  - much HER does not produce recharge 137
  - river flow hydrograph analysis 134–6, 137
  - seasonal variations in baseflow 135, 136–7
- well screens, Drigg boreholes 41
- wetland sediment-Chalk erosional contact 79
- White River Group 64
- winterbournes 100
- Yoredale Shales 193
- zero flux plane (ZFP) technique, for recharge estimation 120–1
- zone of confidence: zone of uncertainty 208