

Index

Note: page numbers in *italic* refer to illustrations; those in bold refer to **tables**

- Accumulation Plateau 240, 241, 242, 243-244, 244
- Adriatic Sea 119
western sector, seismo-stratigraphy surveys 121
- African Planation Surface, Uganda 250
- Ahtme Elevation 75, 77, 77, 78, 81
- Ain Grab Formation 130, 131, 132, 140
- Allan Nunatak 264
- alluvial fans
Guide Dong Shan 190, 190, 196
Nemegt Uul 203, 205, 206, 208, 214, 217
and terraces, Guide Basin, Tibet 190, 190, 192, 196
- Alpine Storm, outer ripples 3
- altiplano, confusion over term 245
- Altyn Tagh Fault 184-185, 184
- Andes 248-250
- Andes, Chilean, fan-like faults 248
- Andes, Ecuadorian 249-251
Cordillera and gravity spreading 249
fold and thrust belt results from Upper Miocene decollements 250
formed by vertical uplift following planation 249, 251
granite/granodiorite batholiths, mainly Mesozoic 250
modelled by single planation surface 245
morphotectonic units 239-246
symmetry round Inter-Andean Depression 250
uplift 246-251
- Andes, Peruvian, block faults 248
- angiosperms 55
- Anglesey 70
- Anglesey–Arfon block 66, 70
- Anglesey–Snowdonian landscape contrast 71
- Anglesey/adjacent Welsh mainland, Menaian Surface/Platform 48-49
- Antarctica, extreme landscape stability 255-267
denudation rate estimates from cosmogenic isotope data 264-265
Dry Valleys region 256-258
rates of landscape change 258-264
- anticlines, crustal ramp 184
- Apaqui River Gorge 240, 242
- apatite fission-tracking studies (AFT) 14-15, 71
indicates major erosional event, eastern flank of Southern Scandes 88
Northern Apennines 121
- Apennine thrust belt, extension and uplift, post-Miocene 109
- Apennines
Northern, Pliocene to present-day uplift and denudation rates 119-125
remnant landsurfaces 110-116
- Appalachian Mountains 250
- Arena Sandstone 265
- Arena Valley
avalanche tongues in 259, 263
clay content of ash 263
- Argentina, palaeo-landsurfaces and neotectonics 229-238
geological setting 229-232
uplift characteristics 235-236
- Asgard Range, glacial landforms 264
- Ashdown Forest/Crowborough–Burwash–Mountfield–Fairlight axis 26, 29
- Asian lithosphere, underlying northeastern margin of Tibetan Plateau 184
- Aterno River, uppermost reach, case study 112-113
- Atlas Mountains 127
- Atlas Thrust Front 128
- Australia, gravity spreading after vertical uplift 249
- avalanche tongues, Arena Valley, retain form on slopes 259, 263
- badlands
Gobi desert 205, 206
Irangi Hills 158
- bajadas *see* alluvial fans
- Bala Fault 66, 67
- Ballyadams Formation 53
- Ballygaddy Outlier, Co. Offaly, saprolite fill 53
- Baltic Oil-Shale Basin, pre-Devonian landscape 75-83
- Baltic Syncline 82
- Barrow Valley, Co. Carlow 50
- Barton Basement Bed, chert in 34-35
- basement
British Isles, a complex mosaic of differing terranes 67
Caledonian, Scandinavia 85
crystalline, Sudetic Foreland 95
Lower Cretaceous, cut by planation surface 239
Precambrian
Baltic Oil-Shale Basin 76
Scandinavia 85
- basement control 13, 28
- basin inversion, south of the Variscan Front 13
- beach gravels, St Agnes sediments 47
- Beacon Cottage Farm Outlier 48
Oligocene sediments 48, 56
and St Agnes Outlier 56
sediments rest on more irregular surface 48, 56
- Beacon Member 56
- Beacon Supergroup 256, 264
- Beacon Valley 263
- bedrock
Baltic Oil-Shale Basin 76-77, 78
deeply weathered, found under Neogene sediments 96-97
Dry Valleys region, intruded by dolerites 256
serpentinite, Niemcza Hills, deeply weathered 100
- bedrock gorges, rectilinear 185
- Bees Nest Pit, Derbyshire, Kenslow Clays in 52
- Beglia Formation 131, 132
- Bekaa Valley
historical evidence for earthquakes 145
Yammouneh Fault forming western boundary 145
- BIRPS, deep seismic-reflection profiling 66
- Black Mountains–Fforest Fawr–Brecon Beacons scarp 66, 68-69
relief inverted with respect to Palaeozoic landscape 69
- Blackdown 25
- Bohemian Massif, Sudetes and Sudetic Foreland 93-105
- Boiano basin case study 113, 115-116
- altitudinal distribution of terrace sequences 115, 115
an inter-montane depression 113
bounded by normal fault-sets 113
SW fault controls tectonic evolution of 115, 116
terrace sequences, anomaly in accordance of 115
- Bosherston–Castlemartin Surface 56
erosional bevel 48
- braid plains, Guide Basin 190, 196
- Braintree Line 35
- 'Breccia di Bisegna' Formation 111
- Brevard Fault Zone, appearance on COCORP profile 69-70
- British Institutions Reflection Profiling Syndicate *see* BIRPS
- British Isles
palaeoforms and saprolites preserved on low coastal platforms 71
sharp lateral discontinuities in continental crust beneath 66
- British margin
receives continuous energy input 71
stress state due to dynamic processes 68
strong control of pre-rifting structure 65-66, 66
- brittle crust 67
in front of Snowdonian front 70
preserves signature of previous tectonic events 68
- bulk-shortening 13
- Bullers Hill Gravel 11
- Bullslaughter Bay syncline, saprolite bodies, chemically weathered mudstone 56

- Caesar's Camp Gravels, reassessed 8
 Calabrian level, ubiquitous nature of pervasive 49-50
 calcrete 208
 Cantalupo sequence 115
 Carboniferous Limestone, Central Ireland, undated saprolitic/palaeosol masses 58
 Cardigan Bay Basin 68
 'Celtic Plain', early Neogene 58
 Central Europe, epirogenic and block uplift, Late Tertiary 93
 Central Range, Taiwan 170, 172
 Cha Lang Valley 196
 braid plain 187, 190, 192, 193, 196
 faults, Guide Formation badlands 193, 194
 Chalk
 complexity of sedimentation and local variation in 30-31
 preservation of outliers indicates long periods of morphostasis 12
 rapid removal of 18
 timing of denudation of 7
 varying dates for exhumation of 40 *see also* Upper Chalk
 Chalklands
 divisible into morphotectonic regions 16
 evolutionary geomorphology, a contemporary synthesis 16-20
 morphostasis, late Palaeogene and Neogene 41
 southern 2-3
 Chambira Formation 245
 Channel High 2
 shows inversion 13
 Channel Uplands, isolation of 16
 Chattian and Miocene deposits, links with 5-150 m surfaces 55-58
 Cherhil Formation 130
 Cherichira Anticline 130, 135, 136, 136, 138
 Cherichira Blind Thrust 133, 135, 136
 Cherichira and Grigema drainage systems, evolution of 138-139
 Cherichira Sole Thrust 135-137
 tectonic uplift during reactivation of 138
 Cherichira Thrust 130, 133-135, 136
 cinder cones, Dry Valleys region, slow erosion of 259
 Clay-with-Flints 7, 11
 climate
 climatic environment, Taiwan 173-175
 Eocene 19, 34
 Gobi Desert 204
 Irangi Hills 158
 present and past, Antarctica 256, 258
 regional and global, affected by the raising of Tibet 183
 climatic cooling, throughout the Oligocene and Miocene 53
 Clogrenan Formation 53
 Coastal Range, Taiwan 169-170, 172
 compression
 and crustal shortening 248
 late Silurian 82
 conifers 55
 continental crust
 anisotropy of 66-8
 imposed stress, brittle and ductile crust, different behaviour 68
 permanent record of geological events affecting it 67
 continental plate boundary system 143
 Cornubian massif 68
 Cornubian terrane 67
 Cornwall, west 51
 wide 50-120 m surface, marine/Pliocene view 45, 47
 cosmogenic isotope data, Dry Valleys region 264-265
 crustal basins, Scandinavia 86
 crustal blocks, morphotectonic equilibrium 67
 crustal discontinuity(ies)
 major, between Snowdon and Anglesey-Arfon blocks 70
 and the origins of major escarpments 68-69
 strongly influence present landscapes 71
 crustal movements, Irangi Hills 162
 crustal thickening 184
 associated with active collision 171, 173
 cuesta topography, extra-glacial Chalklands 2-3
 cuestas, developed on Upper and Lower Greensand 25
 Culver Chalk 31
 Curaray Formation, Late Miocene fauna 245
 Cutucu Uplift 245
 Danghe Nan Shan 184, 184
 Danian 11
 creation of Summit Surface implies rapid removal of Chalk 12
 Dartmoor, as a monadnock 11
 Datong Shan 184
 Dead Sea Transform 143, 145
 and Yammounh Fault, plate tectonic and regional setting 144
 Yammounh Fault not active segment in Lebanon 150
 decollements 250, 251
 deforestation, Irangi Hills 158
 deformation
 compressive 170-171
 Miocene *see* tectonism, Miocene
 Nemegt Uul 215, 217
 neotectonic, poorly understood, southern Argentina 229
 denudation
 Central Weald 38-39, 39
 controlled by bedrock properties 105
 gross, Tertiary, variations, southern England 32-33, 33
 high rates of induced by extreme precipitation 172, 174
 lithologically and structurally controlled 103-104
 net, southern England, since end-Cretaceous 33
 North Downs backslope benches 38, 39-40, 40
 Palaeocene 18
 Palaeogene 32
 Pleistocene 12, 28
 Quaternary, deduced rates of 40-41
 rates in the Northern Apennines 121-122
 removal of Chalk cover, southern England 25
 South Downs crest 38, 39-40, 40
 denudation chronology studies 49-51
 denudation rates, catchments in semi-arid areas 165
 desert pavements, Nemegt Uul 207, 208, 213, 215
 Djebel Cherichira
 aspects of stratigraphy 129, 131-132
 location of 127, 128
 structural uplift 128-129
 structure of thrust front 132-137
 tectonic evolution and structural uplift of thrust front at 132-137
 topographic uplift and erosion of 137-138
 Djebel Cherichira and Oued Grigema, uplift and erosional history of 127-142
 Djebels, Triassic-cored, evolution of in Tunisia 129, 131-132
 Doble Member 56
 Dong Gou Grassland, Guide Basin 181, 187, 188-189, 189
 Dorking-Penshurst-Tonbridge-Biddenden axis 29
 Douhou Louyang river 187, 190
 Dover Straits 30
 discordant, explanations 27
 drainage
 discordant 5, 27
 drainage patterns
 Irangi Hills, modified 161
 southern England, accordant and discordant 7, 9
 Weald, concordant and discordant 27, 27
 Drum Hills, planation staircase system 49
 Dry Valleys region, Antarctica
 active slope processes in coastal zone 258
 climates past and present 256, 258
 denudation rate estimates from cosmogenic isotope data 264-265
 described 256
 evidence demonstrating slow rate of landscape change 260-261
 geological and tectonic setting 256
 glacial landforms 264

- mean denudation rates very low 265
 modification of landforms limited 265
 morphological/depositional
 evidence of rates of landscape change 258-264
 morphology of 256
 persistent low temperature, hyper-aridity and low rates of landscape change 255-256
 progress in geochronometric dating of landforms 255, 258-259
 raised marine features 263-264
 ductile creep 68
 ductile crust, ductility increases with depth 67
 dunes, and desert pavement, Nemegt Uul 206, 209, 213, 215, 217
 dynamic equilibrium 65
- earthquakes
 Aterno River, uppermost reach 112
 Boiano basin 115
 Irangi Hills 161-162, 163, 163
 Lleyn Peninsula 70
 Qinghai Lake and Gonghe Basin 185
- East Africa, suspended-sediment yields from large catchments 166
- East Africa Rift System 157, 158
 data on uplift rates scarce 163
- East Anglia, downwarping of 16, 19
- East Antarctic Ice Sheet 258
- East Devon Plateau
 Combyne Soil remnants 11
 summit mantled by residual soils 11
- Eastern Cordillera (Cordillera Real), Ecuador 241, 242, 243-244, 244-245
 planation surface 245
- Eastern Longitudinal Valley, Taiwan 169, 170, 172
- Ecuador, planation surfaces 239-253
- El Bouqaiia Depression 148, 151
- El Houfia Extensional Faults 130
 history of 137
- end Neogene datum, position of 35-36, 36
- endogenic energy 68
- Eocene
 sub-tropical climate 34
 subaerial erosion in a hot climate 19
- Equilibrium Line Altitude (ELA)
 depressions, Qinghai Nan Shan and Laji Shan 196
- erosion
 assessment of using suspended sediment data 219
 marine
 Cenozoic 58
 Eocene, link with London Basin Chalkland platforms rejected 8-9
 Weald, pulses of in later Palaeogene 34
- erosion cycle 68
- erosional relief, East European Craton, development of 75
- erosional surface, in broad region from Salisbury to Dartmoor 11-12
- escarpments 66, 68-69
 erosional 152
 Jelania Góra basin 102
 origins of and crustal discontinuities 68-69
 separating Irangi Hills from Maasai Plains 157
 straight, associated with high-angle faults 240
 Walbrzych, Middle Sudetes, lithologically controlled 103
 Wesenberg Escarpment 78-79, 82
 Western Cordillera, Ecuador 239
see also Snowdonia front scarp
- Estonia, flatter hilly relief 75
- etchplains, on stable blocks 71
- etchplanation 17-18, 19, 105
 dynamic 65
 widely recognized in the tropics 18
- etchsurfaces
 stripped, Irangi Hills 157
 sub-Cretaceous 87-88
 sub-Jurassic 87-8
 sub-Mesozoic 87, 87-88, 87
 Sudetic Foreland 97-101
 a general view 101
- Eurasian continental plate 169, 171
- Evolutionary model 37, 39, 39
- Faillie du Pas de Calais 27
- fault complexes/systems, strike-slip, left-lateral 184, 202
- fault propagation fold *see* Cherichira Anticline
- faulting
 co-seismic 112
 dip-slip 155
 extensional, Guide Basin 193-194, 196, 198
 Guide Basin 185, 190
 normal 111
 Guide Basin 194
 Xining Basin 194
 Quaternary, Nemegt Uul 215
 strike-slip 170-171
 thrust, and oblique-slip 215
 transcurrent, Plio-Quaternary, evidence against 147-152
- faults 183
 active, central and southern Italy 109
 base Jurassic 26
 Boiano basin 113, 115
 dip at low angles into reflective lower crust 68
 dip-slip normal 109
 divide Irangi Hills into tectonic blocks 160
 extensional
 Ecuador 239, 244
 El Houfia Extensional Faults 130, 148, 151
- fan-like divergence 250
 geometry of at depth, reconstruction 233, 233
 normal, Guide Formation 193, 194
 post-planation 251
 pre-existing, reactivation of in brittle crust 67
 reverse, Sierras Pampeanas 229, 230-231, 232, 235
 sigmoidal curvature 203
 strike-slip 248
see also thrust faults
- Fennoscandian Shield 75, 76
 basic relief types 87
 differing amounts of Neogene and Palaeogene uplift 89
- Ferrar dolerites 256
- Ferrar Glacier 257
- fill terraces 110
- flexural subsidence 138
 regional 128-129
- flexuring
 early Palaeogene 3
 mid-Tertiary 28
- Flimston Outlier/Flimston Pipeclays 48, 56-57
- floods, catastrophic, from cyclones 174
- floras 53, 55
- folds/folding
 and faulting, contemporaneous, indications of 13
 large, Nemegt Uul 204
 Oligo-Miocene 36
 plains-type, Baltic Oil-Shale Basin 77
 small en echelon 28
- foredeep
 Northern Apennines 119-121, 121
- foreland basin development 140
- Fortuna Sandstone Formation 130, 131
- Fucino basin case study 111-112, 112, 114, 116, 119
- Gaohongai River 187, 188
 incision into basin deposits 194, 197-198
- Gash Breccias 48
- geographical information systems 201
- Ghab Fault 145
- gibbsite 57
- glacial landforms, Dry Valleys region 264
- glaciation, Quaternary in northeast Tibet, no firm evidence for 185
- Gobi Altai Mountains 201, 202
 North Gobi Altai 202
- Gobi desert 204-205
- Gobi Platform 184, 185
- Gobi-Tien Shan fault system 201, 202, 202
- goethite 57
- Gonghe Lake Basin 184
- gravels, siliclastic, covering Bosherton-Castlemartin Surface 48
 age of 57
 possibly a pediment 57

- gravity sliding, Ecuadorian Andes 240
- gravity spreading 249, 250
- gravity tectonics, causing compressive folds and thrusts 249
- Great Glen Fault 66, 67, 68, 70
- Grigema Unconformity 130, 131, 135, 136
- Grochowa Massif 95
- Groombridge–Benenden axis 29
- Guajaquil–Babahoyo–S. Domingo Fault 241
- Guayalabamba Gorge, Ecuadorian Andes 242
- Guide Basin, Tibet 185
 - alluvial fans and terraces 190, 190, 192, 196
 - base-level lowering, due to Yellow River drainage 193
 - deposition of molasse in Miocene 185
 - faulting 190
 - flysch deposits thrust and folded 185
 - loess accumulation, Surface 2 194
 - model for development of 197–198
 - molasse deposits suggest closed basin 193
 - reactivation of faults in Cenozoic 185
 - recent small-scale glaciation and deglaciation 195–196
 - Surface 1 192, 193, 197–198
 - surfaces developed in 188–189, 196, 198
- Guide Dong Shan 185, 187, 188, 190
 - high piedmont terrace 190, 190, 193
 - incised alluvial fan 190, 196
 - scarps attributable to faulting 190, 193
 - Surface 1 at foot of 193, 197–198
- Guide Formation, described 185
- Gwna Group 49
- Haiyuan Fault 184
 - activity along 184–185
 - results of changes in trends 185
- Haldon Hills 3, 11
- Hampshire–Dieppe Basin 13, 16, 18
- Hangay Dome 202
- Hardy's multiquadric method 220
- Hastings Beds 41
 - forming 'High' Weald 25
- Haubi basin
 - catchment denudation rate calculated 163–164
 - highly accelerated soil erosion 166
 - increase in apparent sedimentation rate 163
- Haubi, Lake
 - deep sediment profile 159, 161
 - formation of 161
 - severely degraded catchment 158
- Headley Sand 3
 - fossiliferous ironstone problem 8
- high-elevation surfaces, Dry Valleys Region 256
- Hindhead 25
- Hipparion cf. tchicoicum* 185
- Hipparion fassatum* 185
- Hipparion platyodus* 185
- hog's back ridges 2
- Hollymount Outlier, Co. Laois, age of solution-subsidence infill 52–53, 58
- homeomorphism 60
- Homs Basalt 154
 - age of 150–151, 151
 - lies unconformably against Yammouneh Fault 150, 154
 - relationship to Yammouneh Fault 145, 147–150
 - relationships with shattered carbonates 150
- Hythe Beds 25, 36
- Iapetus Suture 66, 67
- ignimbrites, Cuenca, Ecuador 246
- Indian crust, underthrusting Tibetan Plateau 184
- inherited landscapes and uplift, Sudetic Foreland 93–105
- Inter-Andean Depression 240, 241, 242, 243–234, 244
 - Inter-Andean Graben, ophiolites and volcanoes 248
 - Quaternary strato-volcanoes 246
- Intermediate Atlas 128
- intermontane basins
 - central Italy 111
 - compressional 184
- intrusions
 - Andes, Ecuadorian 250
 - Sierras Pampeanas 232
 - Sudetic Foreland 95
 - see also Klodzko-Zloty Stok Granitoid Massif; Strzegom-Sobótka granitoid Massif
- inversion structures see linear inversion structures; upwarping
- inversion tectonics 28–29
 - framework for southern England 34
 - in the Weald 28, 29
- Irangi Hills, Kondoia, Tanzania
 - earthquakes, uplift and natural erosions 161–163
 - geology and climate 157–158
 - N–S fault scarps 157, 159
 - recent sedimentation and denudation rates 163–166
 - tectonics and drainage systems 159–161
 - topographic transects 159, 160
 - ironstone, fossiliferous 8
 - Isle of Wight Monocline 10, 18
 - isostatic readjustment/rebound 65, 123, 174
- Italy, central, palaeolandsurfaces and recent tectonics 109–117
- Izhora Heights 75, 77, 77
- Jabel Akroum ridge 148, 151–152
- Jason glacio-marine diamicton, evidence of 263
- Jelania Góra intramontane basin 98
 - of complex origin 102
 - morphological boundary 102
 - relationships between landforms and bedrock properties 102
- Jumbeli–Naranjal Fault 139
- K–Ar dating, Homs Basalt 150–151, 151, 155
- Kang Zhou Shan, Guide Basin 187, 189–190, 189, 191
- kaolinite, desilication of 57
- Kenslow Clays 52
- Kenslow flora, and evolution of the Pennines 51
- Klodzko region 98
- Klodzko-Zloty Stok Granitoid Massif 103, 104
 - bedrock-controlled topography 103, 104
- Kredija Backthrust 130, 131, 136, 137
- Kredija Strike-Slip Fault 130
 - age of 140
 - truncates El Houfia Extensional Faults and Cherichira Thrust 137
- Laga Formation 119, 120
- Laji Shan 185
- land degradation, Irangi Hills 157
- landform evolution, long-term, neglected 65
- landforms
 - may help to unravel uplift and subsidence histories 93
 - pre-Devonian, Baltic Oil-Shale Basin 77–81
 - regional-scale, controlled by crustal properties 69
 - tectonic, stability of 155
- landscape change, evidence demonstrating slow rate of 260–261
- landscape development, models emphasising time-dependent landforms 71
- landscape differentiation, pre-Neogene, in Europe 105
- landscape evolution
 - part played by deep weathering 96–97
 - reached by Pleistocene ice-sheets 96
 - and solution subsidence 51–53
 - in Southern Mongolia 201–218
 - Sudetes
 - a new look at 102–105
 - and Sudetic Foreland, differing 96
 - Tertiary, new models of 10–13
- landscape stability
 - eastern Taiwan 169–181
 - extreme, Antarctica 255–267
- landscapes, exhumed and partly buried 93
- landslides, and rockfalls, triggered by heavy rain 174
- Latvia
 - extensive tectonic deformation 75

- short hiatus, Silurian to Devonian 82
- Lebanon, landscape evolution in 143-156
 preservation of landscapes 152-153
- Leith Hill 25
- Lenham Beds 3, 8
- Lenham Beds incursion 20
- Lenham surface, controversial 3
- Lewes–Medway line 30
 topography to East more subdued 36
- linear inversion structures/axes/zones 13, 19, 29
 compartmentalize southern England 13-14, 34
- linear zones of disturbance, Baltic Oil-Shale Basin 77
- lithological control
 Chalklands 13
 scarps around Walbrzych, Middle Sudetes 103
- lithosphere
 role of reflective ductile lower crust 68
 typical structure around and beneath the British Isles 67
 typical vertical structure 67-68
- lixisols 163-164
- Llyn Peninsula earthquake 70
- loessic cover, Sierra de San Luis palaeolandsurface 231-232
- London Basin 13
 estimating original thickness of Chalk 31
 platforms on flanks of 17-19, 19
 Upper Chalk border 2
 warping 35
see also Red Crag
- London–Brabant Platform, no sub-horizontal reflections 68
- London–Brabant/Variscan Front, a fundamental boundary 16
- Long Plain Fault, a low-angle thrust 249
- Longyang Gorge 194
- Lower Chalk 2
- lower crust, deforms along ductile shears 68
- Lower Greensand outcrops 35
- lower-Langhian to Messinian interval, two seismic sequences 140
- Luzon Arc 169, 171, 173
- Ma Wu Gorge 188
 formation of 194
- Macchiagodena sequence 115
- Macigno Formation 119, 120
- Mahmoud Formation 131, 132
- Maidstone lineament 29
- Mangan Formation 239
- Manila Trench 169, 171
- mantle intrusion 68
- marine platform, Pliocene 3, 4, 8
- marine shingles, age of changed 11
- marine-cut surfaces, former, criteria for recognition of 50
- Marnoso Arenacea Formation 119, 120
- mass movement, Nemeget Uul 205, 213
- Mejerda Zone 128
- Menai Straits Fault system 66, 67, 70
- Menaian Surface/Platform 48-49, 70
 and the Trwyn y Parc solution-subsidence complex 57
- mesas and buttes, Guide Basin 189, 189, 195
- Metiaoui Formation 130
- mica cooling ages, Larderello geothermal field 121
- Mio–Pliocene Peneplain 3, 5, 6
 concept abandoned 35
 non-existent in type area 7-8
- mobile belts, British Isles involvement with 66-67
- Mogugu River 187, 188
 incision into basin deposits 194, 197-198
 scarp above terrace 190, 191, 195
- Moho 68, 70
- Mongolia, Southern, landscape evolution 201-218
- monoclines 13, 29
 Isle of Wight 10, 18
- morphogenic systems, as closed systems 65
- morphostasis
 Tertiary episodes, southern England 33
 Wealden area 40
- morphotectonic regions 14, 16, 35
- morphotectonic systems 68-71
- morphotectonics 105
 and Cenozoic history, Sudetic Foreland 96
- Mount Lebanon 145
 northern margin 154
 truncated spur on eastern flank of 145, 153
- Mount Lebanon uplift, ancestral boundary 152
- Mount Marine Fault 112, 115
- mountain ranges, orogenic 248-249
- Muddus Plains 88
 correlation with Palaeic surface, Southern Scandes? 89
- mushroom tectonics 249
 Front Range, Colorado 248
- Napo Uplift 245
- Narva–Luga Lowland (Depression) 77-79, 82
- Natural Pits, Hainault, Belgium 52
- Nazza Plate, subduction segment 229, 232
- Nemeget Uul, Southern Mongolia 201
 a broad sigmoidal-shaped restraining bend 202, 203
 catchment areas 205, 210, 211
 flower structure geometry 204, 215
 geographic and structural setting 202, 204
 geographical control of sedimentary facies 217
- geomorphology 204-215
 mountain front sinuosity 205, 208, 215
 possible differential tilting of mountain range 217
 rock types 203, 217
 structural geology 203-204
- Neogene land surface
 height over Central Weald 36
 position and disposition of 33
- Neogene surfaces, extrapolation of away from the London Basin 3, 5
- Neotectonic Map of Italy* 109
- neotectonism, Plio-Pliocene 58
- Netley Heath, bench a flexured facet of Sub-Palaeogene Surface 8
- New Downs Member
 depositional climate 56
 organic material yields Miocene microflora 47
 represents a freshwater ecosystem 56
- Niemcza Hills 94
- etchsurfaces 100-101, 101
- North America, eastern, sediment yields 219-228
- North Downs
 central, soils 8
 Kent, Lenham surface 3
- North Wales, concept of the morphotectonic system 70-71
- North–South Axis, Tunisia 127, 128
 crustal and lithospheric thickness 129
- Northern Scandes 86, 86, 88
- Norwich Crag 16
- Nummulites vascus* marker horizon 131
- oceanic–continental plate collision zone 169
- Ohio River 223, 224, 225
- oil-shale deposits, commercial, Baltic Oil-Shale Basin 76
- Okinawa Trough 169, 171
- oldland–oldland geomorphological relationships, British Isles 60
- ophiolite complexes
 Nemeget Uul 203
 Sudetic Foreland 95
- ophiolite suites, Laji Shan 185
- Ordovician, Baltic Oil-Shale Basin 82
- orogenesis, Andean 229
- orogenies, Precambrian, Ecuador 244-245
- Oued Cherichira 130
 basal Segui Formation
 braided fluvial deposit 132
 debris-flow deposit 138
 facies change 138
 drainage system
 age of 138
 an ephemeral wadi system 127, 129
 antecedent system maintained its base level 137, 138, 140
 not displaced by Kredija Fault 138

- Oued Cherichira (*cont.*)
 time of formation 132
 generation of structural uplift
 137-138, 140
 position of Cherichira Sole Thrust
 137
 time of formation 132, 140
- Oued Grigema 130
 cuts through Ain Grab Formation
 139
 debris-flow fan deposits of Segui
 Formation 132, 139
 fluvial features post-tectonic 139
 similarities to Oued Cherichira 138
 smaller ephemeral wadi system
 128, 129
 younger consequent drainage
 system 132, 140
- outliers
 Chalk 3
 Palaeocene/Eocene 8
see also Ballygaddy Outlier;
 Beacon Cottage Farm Outlier;
 Flimston Outlier/Flimston
 Pipeclays; St Agnes Outlier;
 St Erth Outlier
- overland flow, Hortonian and
 saturated 174
- Palaeocene
 denudation then marine
 encroachment 18
 variation in erosion 32
 Palaeocene–Eocene boundary, affects
 of choice of position 10
- Palaeogene
 distinction between stable areas
 and areas of pulsed uplift 34
 late
 evolution of the Weald during
 33-35
 off-shore record 34
 time of limited erosion 34
 Palaeogene Denudation Model 3-5,
 37, 38
 Palaeogene residuals 3, 8, 32
 Palaeogene sequences, London and
 Hampshire basins 31
 palaeolandscape development 232
 palaeolandsurfaces
 late Palaeogene and Neogene 40
 Miocene, northern Bekaa,
 terminates at an erosional
 escarpment 152
 and neotectonics, Argentina
 229-238
 and recent tectonics, central Italy
 109-117
- Pancanta Fault system, defined by
 secondary faults 235
- Pandivere Heights 75, 77, 77, 78, 81
- Pantano Negro Fault 235
- Pantano Negro Fault shear zone
 235
- partial melting 68
- passive continental margins 65
- Peak Hill Gravel 11
- pediments, Irangi Hills 157-158
- Peleus Till, Wright Valley,
 preservation of upper feather
 edge 264
- Pembroke Peninsula *see* Bosherton–
 Castlemartin Surface
- periclinal
 Cretaceous, becoming Tertiary
 Monoclines 13
 superimposed on Weald–Artois
 Anticline 25
- Pewsey–London Platform inversion
 axis 13
 important structural divide 16-17
- Philippine Sea oceanic plate 169, 171
- phosphorite layer, Baltic Oil-Shale
 Basin 76
- Pinon Formation 239
- Pinon Terrane 248
- Pisayambo Formation 246
- Pizzoli–Barete basin 112, 115
- plains with residual hills, Scandinavia
 87, 88
 Northern Scandes 88
 planed landscape elements 45
- planation 55-58
 double surfaces of *see*
 etchplanation
- planation model, contradicted in
 Germany 105
- planation surfaces
 divide pre-planation from post-
 planation tectonics 250-251
 Ecuador 239-253
 Irangi Hills 160
 mark the end of tectonic regimes
 248-249
 Palaeogene and Miocene 105
 separated by uplift phases, Sudetes
 96
- planed landscape, Palaeogene/
 Neogene age 58
- plate tectonics
 fails to account for growth of the
 Andes 249-250
 and landform development 65
 Taiwan 169-173
see also Tibet
- Plateau Drift 7
- platforms, Chiltern backslopes,
 structurally controlled 8
- Pliocene
 replacement of Northern Alpine
 foredeep 119
 southern Chalklands coastline 3, 5,
 5
- Po Plain–Adriatic Sea basin, volume
 of Holocene sediments deposited
 121
- poljes
 Lebanon 155
- pollen, Oligocene, Beacon Cottage
 Farm 47
- polycyclic surface, early Tertiary 10
- polygonal cracking, Dry Valleys area,
 preservation of volcanic ash in
 259, 262
- Portezuelo Blanco Fault 234
- Porth Swtan, saprolite pocket 49
- Porth Wen, saprolite pocket 49
- Portland–Wight inversion axis 13, 14
- Portland–Wight–St Vallery axis 14
- post-uplift spreading *see* gravity
 spreading
- Potomac River 220, 222, 223, 224,
 225, 226
- precipitation
 Dry Valleys region 256, 258
 high from tropical cyclones 173-174
- pulsed tectonism 7, 10, 12, 18, 28, 34
- pulsed tectonism model *see*
 Palaeogene Denudation Model
- puna surface *see* planation surfaces,
 Ecuador
- Qaidam Basin 184
- Qilian Shan 184, 184, 202
- Qing Shui river 187, 190
- Qinghai Lake 184, 184
- Qinghai Nan Shan 184
- Quartermain Mountains 264
- quartz, *émoinsés luisant* textures 57
- Quaternary
 deduced rates of denudation 40-41
 differential uplift, southern
 England 35-36
 solutional lowering 36
 stability during 353
- Quaternary movements, and Tertiary
 tectonic episodes 16
- rectilinear slopes 259, 265
- Red Crag
 deposition on marine-trimmed
 surface 35
 eastward tilt 16
 known extent of 15
 marine incursion of limited
 geomorphological
 significance 9
 westward extension 9
 importance of 15-16
- redbeds
 Inter-Andean Depression 242
 Sierras Pampeanas 230, 232
- remnant landsurfaces, Apennines
 110-111
- remnant landscape analysis,
 Fucino basin 112
- Reskajeage Surface 48
- planation around the Palaeogene/
 Neogene boundary 56
- Reyueshui River 187
- Rhes-y-cae, Flintshire, solution
 subsidence 51-52
- river/stream capture 152, 153
 Guide Basin 195
- rivers
 Irangi Hills
 ephemeral 160-161
 sandbed 161
 westward-flowing 161
- Roccamandolfi sequence 115
- Ross Embayment 256
- Roum Fault
 probably active transcurrent
 structure 155

- seismicity focused on 145
 Russian Platform 75, 76
 Ryuku Arc 169, 171
 Ryuku Arc tectonic regime 177
 Ryuku collision event 171
 Ryuku Trench 169, 171
- St Agnes Outlier 56, 60
 buried cliff line 47, 56
 deposits
 environment of deposition 56
 fluvialite 47
 possibly coeval to St Erth
 Outlier (Reid) 47
 terrestrial and Miocene 48
 organic material from New Downs
 Member 47
 sub-Miocene unconformity 48,
 56
- St Agnes–Flimston–Trwyn y Parc
 axis, control offered by 58
- St Erth Outlier, Pliocene marine
 fauna in 47
- St Georges Channel Basin 70
- St Ives/Mount's Bay depression 47
- Salisbury Plain Chalklands 16
- Santiago Terrane 248
- Saouaf Formation 130, 131, 132
 dated as Tortonian 131
 tilting of 136
- saprolites 71
 Northern Apennine summits 122
 and solution-subsidence 49, 55-58,
 70
 South Swedish Dome 87
 stripped from Irangi Hills summits
 157
 stripping of, Sudetic Foreland 104
- sarsens 3, 6, 10
- Savana planation theory 58
- Scandinavia, mountains planated
 after Caledonian orogeny 251
- Scandinavian domes, uplift histories
 revealed by 85-91
 sub-Mesozoic etchsurfaces 87-88,
 87
- sea-level
 eustatic fall, Middle Oligocene 34
 lowering of, late Oligocene 55
- seaways, interior Antarctica, early-
 mid Pliocene 258
- secondary structures, growth of in
 mid-Tertiary 5
- sediment yields
 eastern North America 219-228
 world's rivers, and the global
 denudation system 226
- sedimentation
 late Upper Cretaceous, Tertiary
 style 13
 Lower–Middle Pleistocene,
 Northern Apennines 122
 Oligocene, extent unknown 10
- sediments
 Cenozoic 86
 marine
 Plio–Pleistocene, London Basin
 9
- Turonian, at altitude in the
 Sudetes 96
- Miocene, lack of 31
- Oligocene 31, 34, 48, 56
- Palaeogene
 estimating thickness of removed
 sediments 31
 thicknesses used in cross-
 sectional reconstruction 29,
 32-33
 terrigenous, Middle and Upper
 Eocene, lacking beneath
 Channel 34
- Segui Formation 130, 131, 139, 140
 age 131
 back-steepened 136
 braided fluvial deposit 132, 138
 debris-flow deposit 137, 138
 near Oued Grigema watershed
 132, 139
- seismic activity
 high frequency of associated with
 active collision 173
 northeast Tibet 185
- seismic shaking, vulnerability of
 rocks to 173
- Serdj-Ressas Line 128
- shield areas, interpretations of
 denudational history 85
- Sierra de San Luis 229-236
- Sierras Pampeanas
 Devonian intrusions 232
 El Realito–Mesilla del Cura area
 234, 234
 geological setting 229-232
 bounding faults and their
 relationships 230
 Mesilla del Cura block 234
 Neogene uplift 229
 Palaeolandsurface 232-233
 palaeotopographic surface,
 reconstruction of 233, 233
 Rio Nogoli area 234, 235, 235
 uplift characteristics, contribution
 of palaeolandsurface analysis
 to 235-236
- silcrete formation 10
- Sirius Group 264
 marine diatoms in 258
- Sleza Massif(-Radunia) 94, 95
 pre-Neogene etchsurfaces 94,
 99-100, 100
 imperfect correlation bedrock
 structure-topography 99
 septinite slope 99-100
- slope failures, from torrential rain
 174
- slopes, rectilinear 256, 259, 265
- Snowdonia, large topographic feature
 along a major crustal hinge 70
- Snowdonia block 66, 70
- Snowdonia front scarp, differing
 crustal types on either side 70
- soil erosion 174
 importance of climate for 166
 Irangi Hills 157, 162, 164, 166
- soils, inheritance from former
 Palaeogene cover 8
- solution features, density of on the
 Chalk 16, 17
- solution pipes
 pipe fills 3
 Trwyn y Parc 49, 57, 70
- solution subsidence deposits/fills
 51-53, 57, 70
- Songba Gorge 194
- Souar Shale 130
- South American Craton 248
- South Downs, establishment of sub-
 Palaeogene overstep 13
- South Downs inversion axis 14, 29
- South Swedish Dome 86, 86
 and its palaeosurfaces 89
 last rise 87
 post-uplift development of plains
 with residual hills 87
- southern England, Tertiary evolution
 of
 differential uplift 20
 importance of Neogene 36-37
 inversion tectonics framework and
 argument for pulsed
 tectonism 14
 long-term evolution of, important
 developments 13-16
 low relief in Pliocene, inherited 28
 new model of Tertiary landscape
 evolution 18-20
 views of Green/Jones/Small,
 important differences 12-13
- Southern Scandes 86, 86, 88-89
- stepped terrace sequences, Apennines
 110
- Sticklepath-Lustleigh Fault 66
- Sticklepath-Lustleigh Fault Zone
 67-68
- storm surges 173
- strath terraces 110-115
- strato-volcanoes, Quaternary,
 Ecuador 246
- stresses, anisotropic crust, controlled
 by lines of weakness 67, 68
- strike-slip basin reactivation 140
- structural basins, major
 further definition of 18-19
 Jones/Small model 10
- structural compartmentalization 34
 determines morphotectonic regions
 13-14
 individual inversion axes may
 move independently 14
- structural control, direct, lacking in
 Baltic Oil-Shale Basin 82
- structural restlessness, mid-Jurassic
 to end Tertiary 7
- structure–drainage relationship, SE
 England 7
- structures, inherited, control
 geometry of recent structures
 67-68
- Strzegom Hills 94
 boreholes show unevenness of
 weathering front 97
 inselberg-like landscape 97
- Strzegom-Sobótka granitoid Massif
 95

- Strzegom-Sobótka granitoid Massif (*cont.*)
 change in morphology correlates with internal differentiation 97, 99
 etch-surfaces 97, 98, 99
 granite altered to kaolinitic mantle 97
- Strzelin Fault 100
- Sub-Andean Zone and Oriente, Ecuador 245
- sub-Cambrian peneplain
 Scandinavia 85
 South Swedish Dome 87, 87
- sub-Cenomanian surface 60
- Sub-Eocene Surface 3, 4, 6, 11
 tectonic deformation 3
see also Sub-Palaeogene Surface
- sub-Inferior Oolite surface 60
- sub-Liassic surface, Mendips 60
- sub-Mesozoic unconformities 58, 60
- Sub-Miocene facets 10
- sub-Miocene unconformity, beneath St Agnes outlier 48, 56
- Sub-Oligocene facets 10
- Sub-Palaeogene Surface/
 unconformity 3, 17
 deduced position in relation to Chalk 29, 33
 multi-faceted, polygenetic and diachronous 10, 11, 18
 overstep and overlap relations 7, 8, 31-32
- sub-Red Crag surface, planar nature of 16
- sub-Tertiary unconformity 31-32
 fossilized 10
- subduction
 and the Andes 249-250
 at Manila Trench 169
- subsidence, continental 68
- Sudetes 93, 94
 almost total absence of Neogene sediments 96
 basement geology similar to the Foreland 95
 inherited component of landscape 104
 a new look at landscape evolution 102-105
 sedimentary rocks in 95
 uplift
 late Cenozoic 96
 Neogene-Quaternary, and the mountain front 96
- Sudetic Foreland 93, 94, 95-96
- Sudetic Marginal Fault 94, 96
- Summit Plain *see* Summit Surface
- Summit Surface 5
 debate on origin of 28
 evolution under sub-tropical/warm temperate environment 12
 incomplete dissection of 12
 mantled with Clay-with-Flints 3
 survived from early Palaeogene 11
- superficial deposits 3
 contrasting, separation of 11
- tablelands
 developed in Upper Greensand 3
 Devon and Dorset 10-11
- Taiwan
 climatic environment 173-175
 geological setting 169-173
 recent palaeoenvironment 177
 tectonic setting 169, 171
see also Taroko Gorge, Taiwan
- talus deposits, Nemeget Uul 205, 213
- Tanzania, landforms, erosion and deposition 157-168
 earthquakes, uplift and natural erosion 161-163
 geology and climate 157-158
 recent sedimentation and denudation rates 163-166
 studies on erosion and sedimentation in degraded catchments 166
 tectonics and drainage systems 159-161
- Taroko Gorge, Taiwan
 computer modelling of evolution 177-180
 formation of 176-177
 probably on boundary between two tectonic zones 177
 regional setting 175-176
 rock types 176
 sediment yield 176
 stable form along length of 177
 suspended sediment data 176
- Tawonga Fault, a low angle thrust 249
- Taylor Glacier 257
- tectonics
 compressional/compressive 170-171
 contribution of ancient landscape analysis in evaluation of 109-117
 and drainage systems, Irangi Hills 159-161
 Plio-Pleistocene, mountain building 251
see also inversion tectonics
- tectonism
 Cenozoic, northeast Tibet 183
 Cretaceous 31
 in long-term land/landscape development 65
 mid-Tertiary 3, 12, 19, 33
 Miocene 14, 31
 deformation model 27-28, 37, 38
 downgrading of 36-37
 Guide Basin 185, 192
 Neogene 232
 reduced significance of 36
 Palaeogene 18, 31
see also deformation
- Tellian Atlas 127, 128
- Tengouch Lateral Fault 130, 131, 132, 134
 all movement pre-Segui Formation 132
 probably formed synchronously with Cherichira Thrust 132, 134
 reactivation of 135
 uplifted by Cherichira Anticline 135
- Tengouch Thrust 130, 132
- terrace sequences
 Aterno River, uppermost reach 113, 115
 Boiano basin 115
 Fucino basin 112, 114
- terraces
 developed above the Mogogou river 192, 195
 development in Guide Basin may be related to climate change 195
- terranes, exotic, Ecuador, case for 246, 248
- Tertiary
 active stress sources 36
 southern England, evolutionary geomorphology 6
 two periods of exhumation 14-15
 Wooldridge and Linton, emphasis on stability 3
- Tertiary landscape, new models of 10-13
- thrust belts, structural uplift, topographic uplift and erosion in 128-129
- thrust faults 185
 Nemeget Uul 204, 208
- thrust ramp angles, northern and central Tunisia 136
- thrust stacking, causing regional flexural subsidence 128-129
- thrust zone, Nemeget Uul 203
- Tibet 143
 northeast, geomorphology and uplift 183-200
 geological structures 184
 geomorphology south of the Yellow River 188-192
 large-scale gravitational collapse unlikely 194
 tectonic evolution, current opinions 184-185
see also Guide Basin
- Tomoko Hills, probably never deforested 158
- Tornquist Zone 86
- Tower Wood Gravel 11
- Transantarctic Mountains 256
 structural components in the Dry Valleys area 256
- transform-transform-trench triple junction 145
- transgressions
 late Pliocene 8
 no evidence for in Wessex type area 9
 possibility of two 8
 St Erth 60
 Pliocene, extent controversial 19
- Red Crag 9, 15-16, 15, 19, 35
 no morphological evidence outside the London Basin 9
- tree root exposures, Africa, measurements of show erosion rates 164

- Treffynnon Outlier 57
 tropical cyclones (typhoons),
 affecting Taiwan 173
 Trwyn y Parc solution-subsidence
 complex 49, 57
 association with Menaian Surface
 57
 petrography of pipe infills 57
 saprolites
 and organic palaeosols 70
 source differs from host rock 49
 Tunisia *see* Djebel Cherichira and
 Oued Grigema
 Turi Formation 240
 Typhoon Herb 173
 Typhoon Ophelia 174
 typhoons *see* tropical cyclones
 Tyrrhenian margin, volcanic activity
 111
- unconformities
 angular, Sierras Pampeanas 230
 sub-Mesozoic 58, 60
 sub-Miocene 48, 56
 Sub-Palaeogene Surface/
 unconformity 3, 7, 8, 17, 29,
 31-2, 33
 sub-Tertiary 10, 31-2
 see also Grigema Unconformity
 uplift
 assessment of amount and pattern
 105
 and deforestation, causing
 accelerated erosion 164
 differential 35
 and crustal anisotropy 65-74
 Ecuadorian Andes, vertical 249,
 251
 and gully erosion 162-163
 and inherited landscapes, Sudetic
 Foreland 93-105
 and landscape stability, Taroko,
 eastern Taiwan 169-181
 formation of Taroko Gorge 176-177
 late Cenozoic, Sudetes 96
 late Neogene-Quaternary 16
 Mount Lebanon uplift 152
 Neogene 15, 39, 105
 Sierras Pampeanas 229, 232
 Neogene-Quaternary, Sudetes 96
 Northern Apennines, timing of 122
 acceleration in Middle
 Pleistocene 123
 geomorphic impact 122-123
 mode of 122
 uplifted summit areas of low
 relief 122
 permanent, balanced by erosion 68
 Pleistocene, differences as to scale
 and significance of 12
 Plio-Pleistocene, Sierras
 Pampeanas 230
 Plio-Quaternary
 Ecuador 246
 northeast Tibet 183
 Quaternary, Wealden 16
 structural
 Djebel Cherichira 132-137
 Nemegt Uul 203-204, 217-218
 in thrust belts 128-129
 tectonic, Taiwan 171, 173
 Tertiary, Scandinavia 85, 89
 topographic
 Djebel Cherichira 137-138
 in thrust belts 128-129
 transpressional, Gobi Altai 201
 see also upwarping; Weald upwarp
 Upper Chalk
 deposition ceased in Maastrichtian
 18
 estimation of original thickness 30
 outcrop in Chalkland regions 2, 2
 Upper Greensand 3
 upwarping 13, 19, 29
 London Basin Chalk 31
- valley system evolution 151-152, 152,
 153, 154
 Vanda, Lake 263, 264
 Väner Basin 86
 Viivikonna Anticlinical Zone
 structure/topography of the
 sedimentary bedrock 81
 a zone of disturbance 79
 volcanism
 Dry Valleys region
 Cenozoic 256
 volcanic ash deposits 259, 262,
 263, 263
 Eastern and Western Cordillera,
 Ecuador 246
 Inter-Andean Depression 240, 246
 Cuenca area 246
 Mio-Pliocene, southern Argentina
 232
- Wadi Chadra 148
 duration of eruption of Homs
 Basalt into 151
 and Jabel Akroum ridge,
 relationships between
 landscape features 154
 limestone fractured/brecciated east
 of Yammouneh Fault 147-148,
 150
 probably uplifted 155
 section across Yammouneh Fault
 150
 Wadi Charbine 148
 no post-Miocene incision 152
 Wadi ech Chaqif 148
 headwaters captured 152
 Wadi ed Deir 148
 two-stage incision 152
 Wadi Fissane 148
 no post-Miocene incision 152
 Wadi Nsoura 148
 two-stage incision 152
 Wadi Serkhane 148, 151
 headwaters captured 152
 modern valley incised into broader
 valley 151, 152, 153, 154
 Walbrzych, Middle Sudetes 98
 present day morphology 103
 Wali Gong Shan 185
 Walls Boundary Fault 66, 68
- Wardour-Portsmouth inversion axis
 13, 14, 16, 26, 29
 warping
 late prolongation, along Anglesey-
 Arfon and Snowdonia blocks
 hinge 70-71
 London Basin 35
 Palaeogene 36
 and Neogene 14-15
 Quaternary 12
 regional, late Cretaceous 32
 Weald
 background to evolutionary
 geomorphology 25-30
 catastrophist interpretations
 27-28
 fluvial erosional development 28
 inversion tectonics explanation
 28-29
 recognition of anticlinal
 structure 28
 central
 denudation patterns 38-39, 39
 greater uplift 36
 height of Neogene land surface
 over 36
 subaerial denudation possible,
 most of Palaeogene 32
 complex horst structure 16, 36-37
 decoupled from less disturbed
 provinces 29
 established description of 28
 folds, tensional, Alpine
 compressional model
 abandoned 28
 geological reconstruction 30-37
 inversion axes, dominant and
 secondary 29
 models of evolution 37-41
 possible structural partitioning of
 26
 pulsed evolution of 41
 uplift and denudation possible,
 Eocene and Oligocene 34, 35
 variable uplift and later
 downwarping to the East 36
 Weald Island/shoal, London Clay
 Sea, rejected 31
 Weald upwarp 7, 13
 alternative development scenario
 33
 dating of 31
 evidence for early development
 31-32
 Neogene origin? 15
 shown to be asymmetrical in
 reconstruction 30
 shows inversion 13
 Weald-Artois upwarp decoupled
 from more stable areas 41
 well developed in early Palaeogene
 32
 western end experienced greater
 denudation, reasons for 35
 Weald-Artois Anticline 10, 18
 French interpretation of events
 29-30
 removal of Chalk cover 2

- Weald–Artois Anticline (*cont.*)
 uplift and denudation of western
 (Wealden) portion 25-43
- Weald–Boulonnais–Artois horst
 29-30
 possible effect on Weald uplift 32
- weathering
 deep
 of exhumed sub-Cambrian
 peneplain 87
 and pre-Neogene landscape
 evolution 96-97
 grus weathering mantles 102
 weathering residues 3
see also saprolites
- Welsh Basin, northern edge marked
 by Menai Straits Fault system
 70
- Wesenberg Escarpment, Narva–Luga
 Depression 78-79
- Wessex, end-Tertiary surface 36
- western Britain and Ireland
 boreholes penetrating Oligocene
 outliers 53, 55
 broad forested plain at Palaeogene/
 Neogene boundary 60
 Chattian sediments in freshwater
 basins 58
 Chattian/Miocene sedimentary
 outliers and saprolites **54**,
 55-58
 coastal planation surfaces 45-51
- granite areas, elevated since the
 Eocene 105
- oldland fringes represent one of
 oldest little-changed
 landscapes 60
- planar sub-Mesozoic
 unconformities overlain by
 marine sequences 58, 60
- Western Cordillera, Ecuador 239-240,
 241, 242
 bevelled by planation surface 245
- White Chalk
 thickness at outcrop 2
 variability in three dimensions 13
- Wilkes-Pensacola Basin, possible
 seaways across 258
- Wilson cycles 250
- WINCH 4 seismic profile, South Irish
 Sea 70
- winds, low humidity from the Polar
 Plateau 258
- Wooldridge and Linton model 1-2,
 3-5, 37, 38, 39, 49
- Woolwich and Reading Beds,
 overstepping 32
- Wright Valley 263-264
- Wuchia Landslide 174
- Xining (Lake) Basin 184
 normal faulting 194
- Yammouneh Fault 145-148, 150
- follows valley of Wadi Chadra 147,
 149
 geological setting of 143-145
 northern
 drainage evolution, landscape
 correlations across 151-152
 inactive since the Miocene 143
 valleys traced down to Miocene
 palaeosurface, northern
 Bekaa 151-152
- Yammouneh polje 147
 inherited feature 153
- Yellow River
 geomorphology to south of
 188-192
 in the Guide Basin 185, 198
 and the raising of Tibet 183
- Yezhang Grassland, Guide Basin 188,
 189
 deposits beneath 194
 fault scarps delimit extent of 193
 northern edge, scarp with
 triangular facets 190, 192,
 193
- Ying Yao Valley 187, 196
 recent terraces onlap badlands 191,
 192
- Zhihai Shan 185, 187, 192
- Zulova Highland/Massif 94, 95
 inselberg landscape 97, **98**
 kaolin occurrences 97