

# Index

**Note:** Page numbers in **bold** refer to tables; those in *italics* refer to figures (while some are included within page ranges).

- Alberta, Canada, Viking Formation 40, 41, 42, 44,  
45, 49, 50, 53, 55–7, 56
- algal borings 455–6
- Alum Shale Member, North Yorkshire 146, 147, 148
- amalgamated sequence boundaries 51–7
- animal behaviour 399
- ant trace fossils 373, 421, 423  
*see also* Krausichnidae
- Arab-D interval, Saudi Arabia 31
- Archeontomichnus* 434, 439, 446–7
- archosaurs *see* dinosaur tracks
- Arenicolites* 153, 164
- Denison Trough 285, 286, 287
- Ile Formation **241**
- Argentina  
palaeosol ichnofabrics 360–5, 367–71, 375  
Triassic lacustrine deltas 335–54
- Arizona, Bright Angel Shale 213–36
- arthropod trackways 7–8, 313, 318, 321, 322, 326, 346
- Asencio Formation, Uruguay 365–7, 366, 370
- Asteriacites* 164, 165, 166, 167, 168
- Asterosoma* **241**, 249–50, 249, 250, 267, 303
- Attaichnus* 436, 437
- Australia  
Cambro-Ordovician formations 390, 391, 392  
Pebbley Beach Formation 179–212  
shoreface/deltaic facies 273–310
- Austria, Triassic–Jurassic extinction 409–11, 410
- avulsion belt deposits 360, 361
- Bandera Shale Formation, Kansas 162
- barnacle etchings 461–62
- bathymetric reconstruction 4, 72–4
- bayhead delta 55–7, 268
- Beaconites* 317, 321, 365
- bee trace fossils 362, 363, 364, 371, 372, 422, 423  
*see also* Celliformidae
- beetles *see* coleopteran trace fossils
- behavioural homologues 359
- Belichnus* 465
- benthic fauna 146–7
- benthic oxygen levels 398
- Bergaueria* 164, **241**
- BI** *see* bioturbation indices
- Bichordites* 83–4, **84**, 85, 87
- bioerosional trace fossils 455–79  
*see also* borings; etchings
- biogenic marks 224
- biomechanics of dinosaurs 95
- bioturbation  
*see also* ichnofabrics  
claystone and siltstone 281–83  
depth and extent 10–11, 393, 397–8, 408  
intensity 13–14, 183, 296  
laminated sandstone 283–5, 287–8  
measurement 355–8  
muddy sandstone 188–91, 283, 285–7, 286  
mudstone 281  
preserved depth 383  
sandstones 193–4, 193, 283–4  
siltstone 182–8, 279–83  
succession 16–17  
timing 383
- bioturbation indices (BI) 356, 357
- bite traces 466
- bivalve trace fossils 10, 461–62
- Blue Lias, Lyme Regis, Dorset 151–4
- body size 398–99
- borings  
*see also* microborings  
acrothoracicans 464  
algal, cyanobacterial and fungal 457–8  
ascothoracicans 465  
attribution 455–7  
bivalves 461  
Decapoda 465–6  
echinoids 466  
foraminifera 467–8  
gastropods 462–3  
in shells 466–7  
small rosetted 458–9  
sponges 459–60  
worms 460–1
- brachiopod etchings 464
- brackish-water deposits 179–81, 202, 237  
*see also* tidal flats
- Bright Angel Shale, Arizona 213–36  
ecosystem reconstruction 233–4  
facies **220**  
ichnoguilds 223–7  
lithofacies 217–23  
palynofacies 227–30, 228–30  
stratigraphy 215
- bryozoans 455, 463–4
- Burniston dinosaur tracks 113–19, 113, 116, 118
- burrows  
chambered trace fossils 425, 435, 436–8  
dimensions 15  
echinoids and crustaceans 80, 82  
escape 10, 242, 243, 245–6, 246  
Glossifungites ichnofacies 35  
homogenization 11  
horizontal 319  
insects 317, 420  
interstratal 222  
iron-rich sandstones 225  
meniscate 375  
*Phycosiphon* 245

- burrows (*cont.*)  
 Savrda and Bottjer's model 142–3  
 shallow-tier 386, 394  
 thalassinideans 79  
 U-shaped 225  
 vertical 313, 314, 315, 317  
 Bynguano Formation, Australia 391, 392
- Callichurus major* 79, 83  
*Calliopsis* 372
- Cambrian  
 base 399  
 Bright Angel Shale, Arizona 213–36  
 earliest muddy sediments 385–6  
 ichnological record 386–93  
 radiation event 7
- canyons, submarine 37, 38
- Carboniferous tidal environments 157–78
- Cardium Formation, Alberta 49, 51
- Caulostrepsis* 460
- Cavernula* 65, 66–7, 66, 457
- cavities *see* microborings
- Celliclichmus* 362, 363, 364, 423
- Celliforma* 423, 445
- Celliformidae 372, 419, 421, 425
- Centrichmus* 461–62
- chambered trace fossils 419–53  
*see also* insect nests  
 burrows 425  
 construction 420–1  
 fillings 424  
 ichnofamilies 425–44  
 ichnotaxobases 421–25  
 identification **426, 427, 433, 436**  
 shape 421  
 wall types 421–24
- channel environments 261–8, 315–17, 316
- channelized sandstone–mudstone 194–5, 196
- chiton radulation 463
- Chondrites*  
 deep-sea facies 130  
 Denison Trough 282, 288, 304  
 ichnofabric 245, 251, 266, 267  
 ichnoguild 144–5  
 Ile Formation **241**  
 oxygen levels 144, 149–50, 150, 153, 154  
*Chondrites–Skolithos* ichnofabric 242, 242, 266
- Cicatricula* 460
- Circulichnis* 170
- Cladichnus* 131
- claystones 281, 295–7, 360
- climatic control of distribution 77–92
- Clionolithes* 460
- closed lake ichnofaunas 320–1
- coarse-grained facies 274
- coarsening upward sequence 297, 298, 302
- coastal facies 51–2, 194–9, 208, 299
- Cochlichnus* 345, 346, 349
- cocoons 420, 435
- coleopteran trace fossils 364, 365, 371, 372, 421, 422  
*see also* Coprinisphaeridae; Pallichnidae
- colonization styles 16, 17–18
- communities *see* ichnocoenoses; trace fossil assemblages
- composite ichnofabrics 375–6
- Conchotrema* 460
- Conichnus* 289
- continental environments  
*see also* deltaic facies; fluvial facies  
 fluvial ichnofaunas 215–16, 262, 315–19  
 ichnofacies model 312–15, 312  
 lacustrine deltas 335–54  
 lacustrine ichnofaunas 319–23  
 river-dominated facies 253, **344**, 345, 347  
 sequence stratigraphy 323–7
- Coprinisphaera* 312, 368, 369, 425, 427–30, 428, 445
- Coprinisphaeridae 425, 426–32, **427**, 428, 445–6
- Cornichnus* 197
- correlation tool 18–19
- Cretaceous  
 graphoglyptids 134  
 mass extinction event 414–15  
 palaeosols 362–7, 445–6, 447
- cross-bedded sandstones 287, 295  
*see also* hummocky cross-stratification
- crustaceans  
 distribution 81–3  
 ichnology 79–81  
 trace fossils 419, 464–5
- Cruziana* 222, 345, 346, 386, 387  
 preservation 393, 394  
 tidal environments 164, 165, 166
- Cruziana* ichnofacies 7, 41, 42, 45, 49, 284  
 Bright Angel Shale 233  
 diverse proximal expression 301  
 Pebbly Beach Formation 184, 186, 188, 189, 190, 191–3, 197, 208  
 tide-influenced shorelines 161–2
- Cunctichnus* 460
- Curvolithus* 164, 165, 166
- cyanobacterial borings 457–8
- cyclic sediments 199–200
- Cylindricum* 317
- Decapoda, borings 465–6
- deep-sea trace fossils 125–39, 130  
 diversity 125–33, 135  
 time range 126, 127–8
- deltaic facies 237  
 analysis 277–81  
 coarsening-upward sequence 297, 298, 302  
 delta front 252–7, 342–3, 342, 345–6, 346  
 delta plain 343, 343, 348  
 Denison Trough, Australia 273–81, 290–1, 303  
 distributary mouth-bar 253–5, 255, **257**, 258  
 prodeltaic deposits 304, 301

- tide-dominated 237–72, 258  
 Tonto Group 214  
 Triassic rift lakes, Argentina 335–54  
*Dendrina* 459  
*Dendroidichnites* 170  
*Dendrorete* 459  
 Denison Trough Permian sequences, Australia  
   273–310, 275, 276  
   deltaic facies 273–81, 290–301, 303  
   facies analysis 277–81, 282  
   offshore and shoreface deposits 281–90, 306  
 deoxygenation events 142  
*Desmograpton* 131  
 Devonian mass extinction event 400  
*Dictyodora* 128, 129  
*Dimorphichnus* 224  
 dinosaur tracks 93–123, 322  
   anatomy and gait 112  
   experimental work 93–5, 96–100, 107–12  
   laboratory simulations 108–12  
*Diorygma* 468  
*Diplichnites* 170, 171, 345  
*Diplocraterion* 13, 17, 41  
   Denison Trough 284, 286, 285–9  
   ichnofabric 242–3, 242, 266  
   Ile Formation 241  
   oxygen levels 153  
   Pebbley Beach Formation 188, 191, 193, 204, 205  
   tidal environments 164, 165, 167  
   Triassic 403–4, 407  
   wide time range 77  
*Diplopodichnus* 170  
 discontinuities 29–62, 204  
   *see also* omission suites  
   marine flooding surfaces 43, 45, 46, 51–7, 184  
   regressive surfaces of erosion 34–7, 38–44  
   sequence boundaries 34–8, 51–7, 202, 205, 207  
   transgressive surfaces of erosion 39, 43–52, 203,  
   205–7  
 distributary mouth-bar facies 253–5, 257, 258  
 diversity 14–15, 125–33, 135  
   *see also* mass extinction events aftermath  
   bioerosional fossils 468–9  
   changes 78  
   salinity levels and assemblages 172  
   Triassic–Jurassic extinction 408, 413  
 Dolomites, Italy, ichnostratigraphy 403–6  
 Douglas Group, Kansas 162  
 downhole imaging 21  
 drowned river valley 237  
 dung-beetle nests 371, 372, 422  
 duricrusts 365, 367  
 dyads 227, 228, 229  
 dysoxia-trace fossils relationship 141–56  
  
 earthworm trace fossils 373  
*Eatonichnus* 424, 428, 430  
*Echinocardium* 80, 83  
 Echinoid Form 66, 69, 72, 74  
 echinoids 81–3, 466  
 ecology *see* palaeoecology  
 ecosystem reconstruction 233  
 Ekdale and Mason model of oxygenation 144, 150  
*Ellipsoideichnus* 423  
 England  
   Blue Lias, Dorset 151–4  
   Pinhay Bay, Dorset 407–8, 407–8  
   Scalby Formation, Yorkshire 98, 113–19  
   Toarcian, North Yorkshire 145–51  
   Triassic–Jurassic extinction 407–9  
 Entisol 362, 363  
*Entobia* 459–60  
 Eocene 136, 367–71  
 epichnial grooves 224  
 Equisitales 361, 362  
 erosion  
   regressive surfaces 34–7, 38–41, 39, 42, 43  
   transgressive surfaces 39, 43–52, 203, 205–7  
 escape burrow ichnofabric 242, 243, 245–6, 266  
 escape traces 315, 316  
 estuarine facies  
   basin fill 202  
   Bright Angel Shale 213–36  
   channel fill 202, 203  
   differentiation from offshore marine 179–212  
   Ile Formation 252–7, 252, 253  
   palaeoenvironments 265  
   Pebbley Beach Formation 194–9, 208  
   valley fills 53–5  
 etchings 461–3, 464  
 euphotic zone 72  
*Eurygonum* 457  
 experiments on dinosaur tracks 93–5, 96–100,  
   107–12  
 facies 21  
   *see also* deltaic facies; estuarine facies; ichnofacies;  
   lithofacies  
   Bright Angel Shale, Arizona 220  
   characterization 21  
   Denison Trough, Australia 277–81, 282  
   lacustrine deltas 340–5, 341, 344  
   palaeocurrents, Ile Formation, Norway 257  
   southern Sydney Basin 182–99  
   tidal channels 262  
   tide-dominated deltas, Ile Formation 254, 256  
 failure of substrate 104–5, 106–7, 107, 116–19  
 fair-weather conditions 187  
*Fasciculus* 65, 66, 67–8, 72–3, 74, 457  
 feeding traces 314  
 ferricretes 365  
*Fictovichnus* 423, 425, 433–5, 434, 446  
 fillings  
   chambered trace fossils 424  
   estuarine facies 53–5, 202, 203  
 fine-grained sands 109–11

- firmgrounds  
 early Cambrian 386, 393–4  
 Glossifungites ichnofacies 33–4  
 ichnofacies distribution 6  
 lacustrine environments 324, 325, 327  
 omission suites 46, 47, 49
- fish trace fossils 314, 322, 346, 466
- flaser bedding 195, 197
- flashcards 11
- Fleaglellius* 366, 372, 373, 441–2, 440, 446
- flooding *see* marine flooding surfaces
- floodplain ichnocoenoses 315, 319
- fluvial ichnofaunas  
 channels 215–16, 262, 315–17, 316  
 intertidal 169–73, 170, 171  
 overbank 314, 315–17, 318
- flusch trace fossil assemblages 129
- Fontanai* 426–7, 428, 445
- footprints  
*see also* track morphology; trackways  
 dinosaurs 8, 113–19  
 indenter theory equipment 96–7, 98
- foraminifera borings 465–6
- forced-regressive shorefaces 37–43, 39
- Freitag Formation, Australia 299, 300
- freshwater ichnofaunas  
 Bright Angel Shale 213–36  
 fluvial 215–16, 262, 315–19  
 lacustrine 319–23  
 lacustrine deltas 38–47  
 sequence stratigraphy 326–7
- frontiers 19–22
- fugichnia 191, 193, 197
- fungal borings 457–8
- gamma-ray spectrometry 145
- Gastrochaenolites* 462
- gastropod trace fossils 462–3, 467
- genetic stratigraphy 29
- Globodendrina* 66, 69, 71, 459
- Glockerichnus* 131
- Glossifungites ichnofacies 4, 31, 33–4, 35, 49, 56  
 assemblages 36, 38  
 omission suites 41, 42, 46, 47  
 Pebbly Beach Formation 205, 207  
 trace fossil associations 35, 41, 42
- Gordia* 170, 171, 346
- Grand Canyon, Bright Angel Shale 213–36
- graphoglyptids 129–31
- grazing trails 314, 318
- Grebs Nest Point Formation 387, 388–91
- Gyrochorte* 168, 241, 242, 243, 266
- Gyrolithes* 241, 386, 388
- hardgrounds 30, 32
- HCS *see* hummocky cross-stratification
- Helicotaphrichnus* 460
- Helminthoidichnites* 170, 346
- Helminthopsis* 170, 171, 283, 286, 304
- Helminthorhapse* 130
- heterotrophs 72, 73
- hiatus *see* omission suites
- horizontal burrows 319
- hummocky cross-stratification (HCS) 187, 190, 191, 193, 197, 255, 287
- Hyellomorpha* 457
- hypichnial casts 224
- ichnocoenoses 18, 30–1  
 floodplain 315, 319  
 oxygen-related 141–4
- ichnofabrics  
*see also* bioturbation; palaeosol ichnofabrics  
 composite 375–6  
 constituent diagram 12  
 Ile Formation, Norway 239, 241–51  
 marine/non-marine intercalation 20  
 methods of analysis 11–18, 355–8  
 palaeoenvironments 14  
 scales of development 12  
 stacking patterns 18–19, 264, 268–9  
 tidal channels 263
- ichnofacies  
*see also* Cruziana ichnofacies; Glossifungites ichnofacies; Skolithos ichnofacies  
 concept 4–7, 18  
 continental environments model 312–15, 312  
 Corinisphaera 312  
 distribution 30  
 fluvio-lacustrine succession 311–33  
 incised valley complexes 54  
 Mermia 312, 314–15, 316, 319, 320, 322, 324, 335, 348  
 Nereites 4  
 palaeoenvironments 5, 6  
 Pebbly Beach Formation 200, 201  
 Scoyenia 4, 312–14, 318, 320, 322, 324, 326, 327  
 substrate types 30–4  
 Teredolites 32–3  
 Trypanites 32  
 vertebrates 119  
 Zoophycos 4, 41, 42, 45, 144–5
- ichnofamilies *see* insect ichnofamilies
- ichnofaunas  
 fluvial 215–16, 262, 315–19  
 fluvio-estuarine intertidal 169–73, 170, 171  
 intertidal 162–3  
 lacustrine 319–23  
 latest Permian 401–2, 402  
 open-marine intertidal 164–7, 185, 202  
 restricted-bay intertidal 167–9, 167  
 Rhaetian 405, 409–13  
 salinity levels and diversity 172  
 trace fossil assemblages 336–7
- ichnogenera  
 estuarine facies 194–9

- Lyme Regis, Dorset 152–4, 152  
 marine facies 182–94  
 range and diversity 126, 127–8
- ichnogniids  
 Bright Angel Shale, Arizona 223–7  
*Zoophycos–Chondrites* 144–5
- ichnometry 15
- ichnostratigraphy 7–8  
 Cambrian 399–400  
 Coprinisphaeridae 445–6  
 Krausichnidae 446–8  
 Pallichnidae 446  
 Triassic 403–6
- ichnotaxa  
 attribution 359  
 bioerosional 455–79  
 climatic control 78  
 stratigraphic range 77, 128–33, **128**, 129
- ichnotaxobases 421–25
- ichnotaxonomy 3–4  
 chambered trace fossils 419–53  
 composition 72  
 microborings 65–6  
 vertebrate tracks 119–20
- ICZN *see* International Code of Zoological Nomenclature
- IHS *see* inclined heterolithic stratification
- Ile Formation, Norway 237–72  
 conceptual facies model 254, 256  
 facies 255, **257**, 261–8  
 ichnofabrics 241–51  
 palaeoenvironments 252–61, **257**  
 sedimentology 238–41, 240, 251–69  
 trace fossil assemblages **241**
- incised shoreface deposits 37–43, **44**  
 incised submarine canyons 37, 38  
 incised valley complexes 52–5
- inclined heterolithic stratification (IHS) 195, 198, 203
- indenter theory 94–5, 96–7, 98, 102–3
- infaunal communities 15–16, 81–3
- insect ichnofamilies 419–20, 425–44  
 Celliformidae 419, 421, 425  
 Coprinisphaeridae 425, 426–32, **427**, 428  
 Krausichnidae 426, 432, 433–42, **436**, 437, 440  
 Pallichnidae 425, 432–5, **433**, 434  
 stratigraphic range **444**
- insect nests 8, 315, 419, 420  
 ants 421  
 bees 362, 363, 364, 370, 371, 372  
 beetles 364, 365, 371, 372, 420  
 termites 366, 370, 372, 373, 421  
 wall types 421–24, 422
- International Code of Zoological Nomenclature (ICZN) 3–4, 120
- intertidal flats  
*see also* tidal flats  
 fluvial ichnofaunas 169–73, 170, 171  
 model 158–62, 160  
 mud 257–8, 260  
 open-marine 164–7, 164, 166  
 restricted-bay 167–9, 167  
 sand 258–9, 260  
 trace fossil distribution 173
- iron-rich sandstones 219–20, 219, 222–3, 231
- Ischigualasto Formation 360–2, 364, 375
- Ischigualasto–Villa Unión basin 338–40, 339, 340
- Jebel Qatrani Formation, Egypt 367, 370
- Jet Rock Member, North Yorkshire 146, 147
- Joffre Shoreface Complex 49, 50, 55, 56
- Judy Creek Field, Alberta 41, 43, 44
- Jurassic  
*see also* Rhaetian  
 archosaur footprints 8  
 Blue Lias, Dorset 151–4  
 Ile Formation, Norway 237–72  
 mass extinction event 406–14  
 palaeosol ichnofaunas 447  
 Toarcian, North Yorkshire 145–51
- Kansas, USA, outcrops and stratigraphy 158, 159
- Kaybob Field, Alberta 41–3, 42
- key stratigraphic surfaces 19, 29–62
- kinematics of track morphology 103–4
- Konservat-Lagerstätten 398
- Kössen Formation, Austria 410–11
- Kouphichnium* 170
- Krausichnidae 426, 435–43, **437**, 438, 441, 446–7
- Krausichnus* 420, 423, 435–6, 438–9, 438, 443–4
- Kristin Field, offshore Norway 238–41, 238, 240
- laboratory-simulated tracks 108–12
- lacustrine deltas  
 facies associations 340–45, 341, **344**  
 marginal environments 343–5, 343, 347–9  
 trace fossil assemblages **336–7**, 341, **343**, 345–9
- lacustrine ichnofaunas 319–23
- Laguna Palacios Formation 362–5, 363, 364, 375
- lake basin types 324, 325
- laminated mudstone 199
- laminated sandstone 187–93, 284, 285–90
- Lapispecus* 460
- latitudinal distribution 78, 79, 81–3
- Lias *see* Blue Lias; Toarcian
- Lingulella* 213, 233
- Lingulichnus* 168
- liquefaction failure 117–19
- lithofacies, Bright Angel Shale 217–23, **220**
- Lockeia* 164, 165, 166, 167, 168, **241**
- locomotion traces 313, 314, 318, 345, 349
- loess sequence 362, 363
- loosegrounds 30
- Lower Aldebaran Sandstone 292, 297–9, 302
- lower shoreface deposits 192
- lowstand incised shorefaces 37–43, 39, 44
- Lyme Regis, Dorset, Blue Lias 151–4

- Macaronichnus*  
 Denison Trough 289–96, 297–9, 304, 306  
 Pebbly Beach Formation 189, 191, 194
- Maeandropolydora* 461
- Magenta Beds, Arizona 219–20, 219, 222–3, 231
- Mannville Group–Joli Fou Formation contact 51, 52
- marginal lacustrine environment 343–5, 343, 348–9
- marine bioerosion 455–79  
 attribution to trace-maker 455–7  
 borings 457–61, 462, 464–6, 468  
 etchings 461–63, 464  
 radulation 463  
 round holes 466–8  
 stratigraphic range 456
- marine environments  
 climatic control 77–92  
 ichnofabric 20  
 ichnofacies 5–6  
 siliclastic sediments 383–97
- marine facies 253  
 estuarine deposit differentiation 179–212  
 Pebbly Beach Formation 182–94, 202, 208
- marine flooding surfaces (MFS) 43, 45, 46, 51–7, 184
- mass extinction events aftermath 397–416  
 Cretaceous–Tertiary event 414–15  
 late Devonian event 400  
 Ordovician–Silurian event 401  
 Permian–Triassic event 400–406  
 Triassic–Jurassic event 406–14
- Mauretania, microborings 63–76
- maximum zone of deformation (MZD) 100–2
- medium-grained sands 111–12
- meniscate trace fossils 313, 314, 317, 322, 375, 424
- Mermia* 323
- Mermia ichnofacies 312, 314–15, 316, 319, 320, 322, 324, 335, 350
- Mesozoic  
 bioerosion diversity 468  
 graphoglyptids 134  
 trace fossil diversity 125, 126, 128, 131
- methodology of ichnology 9–19
- MFS *see* marine flooding surfaces
- microborings 63–76, 455–7  
 bathymetric interpretation and model 72–4  
 morphology 66–72
- minute round holes 467–8
- Miocene 83–4, 136
- Mirandaichnium* 170
- Missouri, USA, outcrops and stratigraphy 158, 159
- mixed layers *see* bioturbation
- models  
 conceptual facies 254, 256  
 continental ichnofacies 310–13, 312  
 oxygenation 141–4, 150  
 tidal flats 158–62
- moisture/density relationships 105–6, 114–19
- mollusc shell microborings 63–76, 65
- Monesichnus* 430, 431
- Monocraterion* 241
- Monomorphicichnus* 213, 224
- mouth-bar facies 253–5, 253, 255, 265
- mud flats 161
- muddy sandstone 188–91, 281–5
- mudstones 281, 385–6
- MZD *see* maximum zone of deformation
- nektonic fauna 146
- Neogene graphoglyptids 133–6, 134–6
- neoichnology 21, 94
- Nereites* 130, 164, 166, 167
- Nereites* ichnofacies 4
- nests *see* insect nests
- Nevada, USA, Triassic–Jurassic extinction 411–13
- Nododendrina* 459
- nomenclature 3–4, 65–6  
*see also* ichnotaxonomy
- non-deltaic shoreface deposits 281–90
- non-marine ichnofabric 20
- non-marine ichnofacies 5–6
- Norway, Jurassic tide-dominated delta 237–72
- oceanic anoxic event 145
- offshore deposits 281–90  
 lacustrine facies 340–42  
 marine 179–212  
 tide-dominated 261  
 trace fossil assemblages 303, 304
- Oichnus* 466–7
- Oligocene diversity 136
- omission suites 30–4  
 firmgrounds 46, 47, 49  
 Glossifungites ichnofacies 41, 42, 46, 47
- OPB *see* optical pedobarograph
- open lake ichnofaunas 321–23
- open marine ichnofaunas 164–7, 164, 166, 185, 202
- Ophiomorpha* 79, 80–1, 80, 130  
 Cenozoic occurrences 84  
 ichnofabric 244–5, 244, 266
- Ile Formation 241  
 organism distribution 81–3  
 spatangoid trace fossils 83–7
- optical pedobarograph (OPB) 97–9, 112
- ORB *see* oxygen-restricted biofacies
- Ordovician  
 Grebs Nest Point Formation 387, 388–91  
 ichnological record 386–93  
 mass extinction event 400, 401
- Orthogonum* 65, 66, 68–70, 69, 74, 458
- Ostrebium* 72
- overbank ichnofaunas 317–19, 318
- oxygen-restricted biofacies (ORB) 145, 146–8, 151–2
- oxygenation 8–9  
 evidence in Toarcian deposits 146–8  
 lacustrine environments 314, 322  
 Lias, Dorset 151–2

- oxygenation (*cont.*)  
 measurement 145  
 models 141–4
- Pacoota Sandstone Formation, Australia 15, 391
- palaeobathymetry 4, 72–4
- palaeoclimate criteria 78
- Palaeoconchelis* 73, 458
- palaeoecology  
 analysis 398–9  
 Bright Angel Shale, Arizona 213–36  
 microborings 63–76  
 Tonto Group 214–16
- palaeoenvironments  
 analysis 1, 12, 395–6, 414–15  
 delta front 252–7  
 estuarine 265  
 ichnofabrics 14  
 ichnofacies 5, 6  
 Ile Formation, Norway 257  
 tidal channel 261–8  
 tidal flats 173, 257–61, 260  
 wave- and tide-influenced 259, 290–2
- Palaeogene diversity 126, 127
- palaeogeography, eastern Grand Canyon 216
- Palaeophycus* 221, 222, 346  
 Denison Trough 282, 284, 288–90, 293, 304  
 ichnofabric 242, 243, 248, 266, 267  
 Ile Formation 241  
 oxygen levels 153  
 Pebbly Beach Formation 185, 187, 189  
 tidal environments 164, 167, 168
- Palaeosabella* 461
- palaeosol ichnofabrics 355–82  
 analysis 355–8, 371–2  
 Argentina 360–5, 367–71, 375  
 Asencio Formation, Uruguay 365–7, 366, 370  
 bedding and structures 356–7  
 composite ichnofabrics 375–6  
 Jebel Qatrani Formation, Egypt 367, 370
- palaeosols  
 chambered trace fossils 419–53  
 ichnotaxa range 436
- palaeotopography 162, 214, 216, 216
- Palaeozoic  
 arthropod trace fossils 7–8  
 bioerosion 468  
 diversity 125, 126, 127–31  
 graphoglyptids 133–4  
 ichnofabrics 383–96
- Pallichnidae 425, 432–5, 433, 446
- Pallichnus* 423, 425, 434, 435, 446
- Palmiraichus* 422, 423
- palynofacies 213–14, 227–30, 228–30
- palynomorphs 231–3
- Parowanichnus* 434, 437
- pascichnia traces 150
- Pebbly Beach Formation, Australia 179–212  
 discontinuity surfaces 204  
 estuarine facies 194–9  
 ichnofacies 200, 201  
 marine facies 182–94  
 sequence stratigraphy 199–207, 200, 201, 206  
 trace fossil summary 202  
 transgressive surfaces of erosion 203, 205–7  
 pedofabric/ichnofabric ternary diagram (PITD) 357, 358  
 pedofabrics 357, 360, 368, 371–2  
 pellets 79  
 Pennsylvanian stratigraphy, USA 159  
 permeability of substrate 106–7  
 Permian  
 Denison Trough, Australia 273–309  
 ichnofaunas 401–2, 402  
 mass extinction event 400–6  
 Pebbly Beach Formation, Australia 179–212  
 petroleum industry 19–21  
*Petroxestes* 462  
*Phoebichnus* 241, 245, 246, 266  
 photoautotrophic endoliths 72  
*Phrixichnus* 468  
*Phycodes* 223, 388  
*Phycosiphon* 130  
 burrows 245  
 Denison Trough 282, 281–3, 288, 287, 292–6, 297–301, 306  
 ichnofabric 245, 246, 247, 251, 266  
 Ile Formation 241  
 Pebbly Beach Formation 185, 187, 188, 191, 194  
 Pinhay Bay, Dorset, England 407–8, 407–8  
 piped zones 142, 143  
 PITD *see* pedofabric/ichnofabric ternary diagram  
*Planobola* 457, 467  
*Planolites* 80, 153  
 Denison Trough 283, 287, 293  
 ichnofabric 245–9, 266, 267  
 Ile Formation 241  
 Pebbly Beach Formation 185, 187, 188, 195, 197  
*Planovolites* 468  
 plant fossils 367  
*Platydendrina* 459  
*Plectonema* 72  
 Pleistocene assemblages 85, 86–7  
 Pliocene assemblages 84–5, 86  
*Podichnus* 464  
*Polyactina* 65, 66, 68, 458  
 porous plate box 96–7  
 preservation  
 controls, track morphology 93–123  
*Cruziana* and *Rusophycus* 393, 394  
 potential of lacustrine ichnofaunas 327  
 quality 384–5  
 styles 385  
 prodeltaic deposits 304, 305  
 progradational cycles 274–7  
 prograding shoreface succession 300

- Protovirgularia* 164, 165, 166, 167, 168  
*Psammichnites* 164, 166, 167, 168, 197  
 Pygmy Form 65, 66, 71, 74  
 pyrite framboids 145, 151–2
- quantification of bioturbation 10–11, 355–7
- radiation events 7  
 radulation 463  
*Radulichnus* 463  
*Ramodendrina* 459  
 Ravenscar, Yorkshire, Toarcian deposits 145–51  
 ravinement 43–5, 49  
*Rebuffoichnus* 364, 365, 372, 432, 434, 445  
 Recent microborings 63–76  
 red-bed succession 365–7  
 regressive surfaces of erosion (RSE) 34–7, 38–41, 39, 42, 43  
*Renichnus* 462–3  
 repartition 349  
 research landmarks 3–9  
 reservoirs 21, 273  
 residual strength of substrate 106–7  
 restricted-bay intertidal ichnofaunas 167–9, 167  
*Reticulina* 66, 69, 70, 458  
 Rhaetian ichnofaunas 405, 409–14  
*Rhizocorallium* 149, 153, 164  
   Denison Trough 284, 286, 288, 287, 289  
   Ile Formation **241**  
   Pebbley Beach Formation 185, 188, 191  
   Servino Formation 405, 406, 412  
*Rhopalia* 458  
 rift basins 274  
 rift lakes 335–54  
 ripple structures 195, 196, 243, 253  
 river-dominated facies 253, **344**, 345, 347  
   *see also* fluvial ichnofaunas  
*Rogerella* 464  
*Rosellichnus* 423, 424  
*Rosselia*  
   Denison Trough 278, 284, 288, 287, 292, 296, 300  
   ichnofabric 249–50, 249, 267  
   Ile Formation **241**  
   Pebbley Beach Formation 185, 187, 188, 191, 194  
*Rotundusichnium* 130  
 RSE *see* regressive surfaces of erosion  
*Rusophycus* 166, 386, 388, 391, 392, 393, 394
- Saccomorpha* 65, 66, 67, 74, 456  
 salinity levels 81–2, 172  
 sandbars 160, 231  
 sands 109–12, 258–9, 260  
 sandstones  
   bioturbated 193–4, 193  
   cross-bedded 290, 295  
   deltaic 290–301  
   laminated 187–93, 284, 285–9  
   muddy 188–91, 286, 284–7, 288  
   sheet-like 195–9, 197  
   silty 281–4, 295–7  
   tabular 291–5  
   tidal channels 17  
 sandy siltstones 185–7, 186  
 Sarmiento Formation, Argentina 358, 367–71  
 Savrda and Bottjer model of oxygenation 141–4  
 SB *see* sequence boundaries  
 Scalby Formation, Yorkshire 98, 113–19  
 scale of observation 10  
*Scaphichnium* 425, 433, 434, 446  
 scarabaeid beetles 435  
 scars 463  
*Schaubcylindrichnus* **241**, 243, 244, 247, 248, 266, 267  
*Scolectia* 66, 69, 70–1, 72, 458  
*Scolicia* 81, 82, 83, **84**, 85, 130  
 Scoyenia ichnofacies **4**, 312–14, 313, 318, 320, 322, 324, 326, 327  
 sediment feeders 223  
 sedimentary structures 163, 164, 276  
   *see also* cross-bedded sandstones; soft sediment deformation  
 sedimentology  
   context 10  
   ichnofabrics 4  
   Ile Formation, Norway 239–41, 240, 251–69  
   intertidal flats and subtidal sandbars 160  
   lacustrine deltas 347  
   Pebbley Beach Formation, Australia 179–212  
 Seilacherian ichnofacies 4–7, 18  
 semi-quantitative methods 355–6  
 sequence boundaries (SB) 34–7, 38, 51–7, 202, 205, 207  
 sequence stratigraphy 9  
   case studies 34–51  
   fluvial ichnofacies 326–7  
   lacustrine ichnofacies 324, 325–6  
   Pebbley Beach Formation 199–207, 200, 201, 206  
 Servino Formation, Italy 403–6, 405  
 shallow marine siliclastics 383–96  
 shallow-tier burrows 386, 394  
 shear surfaces 107  
 sheet-like sandstone–mudstone 195–9, 197  
 shelf storm deposits 230–1  
 shell borings 465, 466–7  
 shorefaces  
   comparison with delta front 306  
   facies **278**, 281–90  
   forced regressive and lowstand incised 37–43, **44**  
   prograding 300  
   transgressively incised 39, 47–51, 50  
 siltstones 182–8, 184, 281–4, 297  
 silty sandstones 281–3, 295–7  
 Silurian mass extinction event 400, 401  
 simulated dinosaur tracks 98, 108–12, 109, 110  
*Siphonichnus* **241**, 250–1, 250, 267  
*Skolithos* 315  
   Denison Trough 287, 288, 289



- ichnofabric 247, 248, 267, 285, 289  
 Ile Formation **241**  
 palaeosol 365  
 Pebbly Beach Formation 194, 195, 197  
 piperock 391  
 Skolithos ichnofacies **4**, 42, 53, 161, 188, 312, 316, 320, 321, 324, 327  
 Skolithos–Cruziana ichnofacies 188, 195, 208, 284  
 small borings 458–9, 467–8  
 soft sediment deformation structures 293, 293, 304  
 softgrounds 6, 30, 41, 42, 324, 325, 327  
 soil mechanics 105  
 soil moisture content 105–6, 114–19  
 soils *see* palaeosols  
*Solemya* 166  
 soupgrounds 30  
 spatangoid trace fossils 80, 81, 83–7  
*Spirichnus* 461  
 sponge borings 459–60  
*Spongeliomorpha* 80–1  
 stacking patterns 18–19, 264, 268–9  
*Stiallia* 170, 171  
*Stiaria* 170, 171, 345  
 stiffgrounds 30  
 storm deposits 187–8, 189–91, 230–1, 287–8, 385  
 stratigraphic range 77  
   bioerosive ichnotaxa 456  
   ichnotaxa 77, 128–33, **128**, 129  
   insect ichnotaxa **436**  
   lacustrine delta assemblages **336–7**  
 stratigraphic surfaces 19, 29–62  
 stratigraphy *see* ichnostratigraphy; sequence  
   stratigraphy  
 striated traces 314  
 sub-recent microborings 63–76  
 subaerially exposed surfaces 51–2  
 submarine canyon incision 37, 38  
 substrate-controlled ichnofacies 30–4  
 subsurface tracks 99, 104–5, 109–12  
 subtidal environments 158–62, 160, 259–61, 260  
 subtropical associations 83–5  
 supratidal zone 160–1  
 surface traces 225, 226  
 surface track simulations 99, 108–9  
 surfaces  
   *see also* discontinuities  
   key stratigraphic surfaces 19, 29–62  
   shear surfaces 107  
   subaerially exposed 51–2  
 suspension feeders 223–5  
 Sydney Basin, Australia 179–212  
*Syntermesichnus* 440, 442, 446  
  
 tabular sandstone 288–92  
*Tacuruichnus* 437, 439, 446  
*Taenidium* 153, 185, 315, 362, 363, 365, 375  
   ichnofabric 244–5, 244, 248–9, 249, 266, 267  
   Ile Formation **241**  
  
*Talpina* 461  
 taphonomic pathways 316, 320  
 taxonomy *see* ichnotaxonomy  
*Teichichnus* 167, 168, 389, 390  
   Bright Angel Shale 223  
   Denison Trough 281, 285, 303  
   Ile Formation **241**  
   Pebbly Beach Formation 185, 187, 189–91  
*Teisseirei* 366, 372, 421, 424, 430–2, 431, 445  
 temperate association 85–7  
*Teredolites* 462  
 Teredolites ichnofacies 32–3  
 termite nests 366, 370, 372, 373, 421, 422–3  
   *see also* Krausichnidae  
*Termitichnus* 441–2, 440, 443–4, 446  
 Tertiary  
   diversity 136  
   graphoglyptids 133–6, 134–6  
   insects 8  
   mass extinction event 414–15  
   palaeosol ichnofaunas 444–6  
 tetrads 227, 228, 229  
 Tetrapod tracks 170  
 textures *see* ichnofabrics  
 thalassinidean crustaceans 79, 81  
*Thalassinoides* 41, 80–1, 86, 87, 153  
   ichnofabric 167, 248, 249, 250, 251  
   Ile Formation **241**  
   Triassic 406, 412  
 tidal channel environments 17, 253–6, **257**, 261–8  
 tidal flats 157–78  
   *see also* subtidal environments; supratidal zone  
   facies associations 260  
   ichnofaunas variability 163–73  
   model 158–62, 160  
   palaeoenvironments 257–61, 260  
   sedimentary structures 163, 164  
   stratigraphic implications 173–4  
   successions 259  
   variability 175–8  
 tide-dominated deltas 237–72, 259  
 tiering 15–16, 78, 82  
   Blue Lias 153  
   diagrams 357–8, 397  
   oxygenation events 141–4, 142  
   palaeosols 357, 358, 362, 363, 376  
 time range and diversity 126, 127–8  
 Toarcian deposits, North Yorkshire 145–51  
*Tonganoxichnus* 170, 171  
 Tonto Group, palaeoecology 214–16  
*Topsentopsis* 460  
 trace fossil assemblages  
   *see also* ichnofacies  
   Blue Lias, Dorset 152–4  
   continental 311–33  
   deep-sea ichnotaxa 128–33  
   delta front 345–6, 346  
   delta plain 348

- trace fossil assemblages (*cont.*)  
 distribution controls 349–50  
 diversity and salinity levels 172  
 dyoxia relationship 141–56  
 graphoglyptid proportions 132, 133  
 Ile Formation, Norway 241  
 intertidal 173  
 lacustrine deltas 336–7, 341, 345–9, 346, 347  
 lacustrine sequence stratigraphy 324  
 mass extinction events aftermath 397–418  
 Mesozoic 131  
 Neogene 131  
 palaeosols 359–60  
 Palaeozoic 129–31  
 Pebbly Beach Formation 202  
 prodeltaic v. offshore deposits 303, 304  
 proximal Cruziana ichnofacies 191, 192  
 repartition 349  
 Toarcian, North Yorkshire 147, 148, 149–50
- track morphology 93–123  
 Burniston dinosaur tracks 113–19  
 description and terminology 100–2  
 experimental work 93–5, 96–100, 107–12  
 indenter theory 94–5, 96–7, 102–3  
 moisture/density relationships 105–6, 114–19  
 surface and subsurface features 99–100, 101
- trackways  
 arthropods 7–8, 313, 318, 321, 322, 326, 346  
 vertebrates 313, 318, 322
- transgressive surfaces of erosion (TSE) 39, 43–52, 203, 205–7
- transgressively incised shorefaces 39, 47–51, 50  
*Treptichnus* 170, 171, 346, 386, 388, 399
- Triassic  
 archosaur footprints 8  
 ichnostratigraphy 403–4  
 lacustrine deltas, Argentina 335–54  
 mass extinction events 399–414  
 palaeosols 360–2, 446–7
- Triassic–Jurassic mass extinction event 406–16  
 Central Austria 409–11, 410  
 England 407–8, 407–9  
 Nevada, USA 411–13
- Trichichnus* ichnofabric 251, 267  
*Trichophycus* 387, 389–90
- tridactyl dinosaur tracks 98, 113, 116, 118
- Tripartum Form 66, 69, 71, 74
- trophic reconstruction 231–3
- tropical associations 83–5  
*Trypanites* 461  
 Trypanites ichnofacies 32  
 TSE *see* transgressive surfaces of erosion
- tsunami beds 414–15
- tubular trace fossils 374–5
- tuffaceous palaeosols 362–5, 363, 368
- tunnels 67–8  
*see also* burrows; graphoglyptids
- turbidites 322, 325–6
- U-shaped burrows 225
- UK *see* England
- Ultisols 366, 370  
*Undichna* 170, 171, 345, 346
- upwelling areas 63–76
- Uruguay, Asencio Formation 365–7, 365, 370
- valleys 52–5, 162, 237
- vertebrates  
 body fossils 367  
 dinosaur tracks 93–123  
 ichnofacies 119  
 track taxonomy 119–20  
 trackways 313, 318, 322
- vertical burrows 313, 314, 315, 317
- Vertisols 360–2, 361, 364
- Viking Formation, western Canada 40, 41, 42, 44, 45, 49, 50, 53, 55–7, 56
- Vondrichnus* 440, 441, 446
- walls in chambered trace fossils 421–24
- wave-dominated environments 259, 291, 344, 345–6, 347
- wavy bedding 195, 196, 197
- woodgrounds 30, 32–3
- worm borings 460–1
- wrinkle marks 226
- Yorkshire, UK  
 Burniston dinosaur tracks 113–19, 113, 116, 118  
 Toarcian 145–51, 146
- zone fossil characteristics 7  
*Zoophycos* 284, 286, 287, 306, 406  
*Zoophycos* ichnofacies 4, 41, 42, 45, 144–5  
*Zoophycos–Chondrites* ichnoguuild 144–5