

Contents

| | |
|---|-----|
| PEDLEY, M. & CARANNANTE, G. Cool-water carbonate ramps: a review | 1 |
| NALIN, R., BASSO, D. & MASSARI, F. Pleistocene coralline algal build-ups (<i>coralligène de plateau</i>) and associated bioclastic deposits in the sedimentary cover of Cutro marine terrace (Calabria, southern Italy) | 11 |
| BASSO, D., MORBIOLI, C. & CORSELLI, C. Rhodolith facies evolution and burial as a response to Holocene transgression at the Pontian Islands shelf break | 23 |
| BASSI, D., CARANNANTE, G., MURRU, M., SIMONE, L. & TOSCANO, F. Rhodalgae/bryomol assemblages in temperate-type carbonate, channelized depositional systems: the Early Miocene of the Sarcidano area (Sardinia, Italy) | 35 |
| KERSHAW, S. & GUO, L. Pleistocene calcified cyanobacterial mounds, Perachora Peninsula, central Greece: a controversy of growth and history | 53 |
| FORNÓS, J.J. & AHR, W.M. Present-day temperate carbonate sedimentation on the Balearic Platform, western Mediterranean: compositional and textural variation along a low-energy isolated ramp | 71 |
| TOSCANO, F., VIGLIOTTI, M. & SIMONE, L. Variety of coralline algal deposits (rhodalgae facies) from the Bays of Naples and Pozzuoli (northern Tyrrhenian Sea, Italy) | 85 |
| MASSARI, F. & CHIOCCI, F. Biocalcarene and mixed cool-water prograding bodies of the Mediterranean Pliocene and Pleistocene: architecture, depositional setting and forcing factors | 95 |
| BRAGA, J.C., MARTÍN, J.M., BETZLER, C. & AGUIRRE, J. Models of temperate carbonate deposition in Neogene basins in SE Spain: a synthesis | 121 |
| PEDLEY, M. & GRASSO, M. The response of cool-water carbonates to eustatic change in microtidal, Mediterranean Quaternary settings of Sicily | 137 |
| REUTER, M., BRACHERT, T.C. & KROEGER, K.F. Shallow-marine carbonates of the tropical–temperate transition zone: effects of hinterland climate and basin physiography (late Miocene, Crete, Greece) | 157 |
| RUBERTI, D., TOSCANO, F., CARANNANTE, G. & SIMONE, L. Rudist lithosomes related to current pathways in Upper Cretaceous temperate-type, inner shelves: a case study from the Cilento area, southern Italy | 179 |
| HALFAR, J., STRASSER, M., RIEGL, B. & GODINEZ-ORTA, L. Oceanography, sedimentology and acoustic mapping of a bryomol carbonate factory in the northern Gulf of California, Mexico | 197 |
| LUKASIK, J. & JAMES, N.P. Carbonate sedimentation, climate change and stratigraphic completeness on a Miocene cool-water epeiric ramp, Murray Basin, South Australia | 217 |
| ANASTAS, A.S., DALRYMPLE, R.W., JAMES, N.P. & NELSON, C.S. Lithofacies and dynamics of a cool-water carbonate seaway: mid-Tertiary, Te Kuiti Group, New Zealand | 245 |
| KINDLER, P., RUCHONNET, C. & WHITE, T. The Southern Marion Platform (Marion Plateau, NE Australia) during the early Pliocene: a lowstand-producing, temperate-water carbonate factory | 269 |

| | |
|---|-----|
| HENDY, A.J.W., KAMP, P.J.J. & VONK, A.J. Cool-water shell bed taphofacies from Miocene–Pliocene shelf sequences in New Zealand: utility of taphofacies in sequence stratigraphic analysis | 283 |
| MUTTI, M., JOHN, C.M. & KNOERICH, A.C. Chemostratigraphy in Miocene heterozoan carbonate settings: applications, limitations and perspectives | 307 |
| KNOERICH, A.C. & MUTTI, M. Epitaxial calcite cements in Earth history: a cooler-water phenomenon during aragonite-sea times? | 323 |
| CARON, V., NELSON, C.S. & KAMP, P.J.J. Microstratigraphy of calcite cements in Pliocene cool-water limestones, New Zealand: relationship to sea-level, burial and exhumation events | 337 |
| Index | 367 |