

# Index

References to figures appear in *italic type*

References to tables appear in **bold type**

All entries refer to the Lewisian unless otherwise noted

Abloviak shear zone, Labrador 253, 254, 255, 256, 256  
    compared boundary zone, W. Greenland 257, **257**  
accretion, Scourian 7, 8–9, 11  
accretion model, Napier complex, Antarctica, near  
    isobaric cooling 290, 293–4  
Achiltibuie area  
    garnet-pyroxene thermometry 83, 84  
    two-pyroxene thermometry 83, **84**  
agmatite complex, Gruinard Bay 14, 57, 59, 81  
Akilia supracrustal association, W. Greenland 261, **261**  
    compared E. Hebei Province 303–4  
Albany–Fraser mobile belt 277, 278, 281–3, 282  
Ameralik dykes, W. Greenland  
    emplacement of 217, 247, 261, **261**  
    feeders of Malene metavolcanics 261–2, 271  
    geochemical difference 264, 265, 266, **268**, 271–2  
Amitsøq gneisses, W. Greenland 217, 236, 247, 248, 261, **261**, 262  
    *see also* Ameralik dykes, Fiskenaasset complex, Malene metavolcanics  
amphibolite facies metamorphism  
    age of 16–17, 46  
    cause of 16  
    distribution 14–15  
    dykes 8, 205–6, 227, 231  
    examples of 18  
        Assynt area 168  
        Gruinard Bay 14, 58, 59, 81–2  
        Outer Hebrides 36, 41, 43, 175, 175, 176, 181  
        Scourie area 75  
        S. Mainland 140, 147, 153, 207  
        Stoer area 94  
    magnetic and seismic surveys 187, 188, 197  
    pressure–temperature conditions 36, 88–9, 89, 90, 94, 95, **95**  
    Antarctica **289**, 291–2, 292  
    Greenland 236, 237, 240, 242, 243, 248, 250  
    Labrador 253, 254  
    W. Australia 281, 282, 283  
amphibolites  
    Gruinard Bay 57, 58, 59  
        geochemistry 61, **62**, **63**, 69  
        partial melting 67, 68–70, 69  
        petrogenesis 63–4, 69  
        petrography 60  
        rare earth elements 61, 63, 64, 65, 69  
    Outer Hebrides 30  
    Scourie area 81  
    Stoer area  
        partial melting conditions 97, 101  
        petrology **94**, 102  
    Torridon shear zone 154

E. Hebei Province, China 298, 299, 300, 301, 303  
E. Greenland 238  
    *see also* Malene metavolcanics  
    Labrador 255  
Andean model, explanation of—  
    metamorphic conditions 46  
    Scourian accretion 8, 9  
Angmagssalik charnockite complex 236, 242  
anorthosite complexes  
    formation of 47, 50  
    Greenland and Outer Hebrides compared 8, 244  
    Labrador 254  
Antarctica, East, petrogenesis of high grade terrain 48–9, 53  
    *see also* Napier complex  
aplite veins, Outer Hebrides 37  
apparent polar wander, use of palaeomagnetic record 205  
Archaean  
    Proterozoic boundary, change in magma production 11, 54  
    terrains compared 22–3, 257, 257  
    *see also* Nain Province, Pilbara craton, Scourian, Yilgarn Block  
Ashburton Trough, Australia 277, 278  
Armada marble 109, *110*  
Assynt granulite zone  
    metamorphic conditions 89, 89–90, 165, 168  
    Scourie dykes in 218  
Atlantic, opening of 10, *136*, 137  
augite-rich dyke margins 223, 227  
  
Badcall dyke, remanence 206  
Badcallian granulite facies metamorphism  
    age of 15, 16, 17, 46–7  
    causes of 17, 73  
        deep derived fluids 74  
        enhanced mantle heat flow 74–5  
        tectonic thickening 75–8, 77, 77  
        magmatically enhanced heat flow 73–4  
    equilibrium conditions  
        feldspar thermometry 87–8, 87  
        garnet-pyroxene thermometry 83–5, 84, **85**  
        iron-titanium oxide 85–7  
        previous work 82, **82**  
        Stoer area 94–6, 95, **95**  
        two-pyroxene thermometry 83, **84**  
    evidence from  
        Assynt area 89, 90, 165, 168  
        Gruinard Bay 14, 58, 59, 59, 60, 81  
        magnetic and seismic surveys 187, 188, *191*  
        Outer Hebrides 29  
        Outer Hebrides Fault Zone 175, 175, 181, 187  
        Scourie area 59, 59, 81–2  
        Stoer area 93, 94, 96  
        retrogression 88–90, 89

- banded iron formation  
 S. India 303  
*see also* E. Hebei Province
- Bangemall Basin 277, 278
- Beannach dyke, transverse modal variations and margins 219–20, **219**, 220, 228
- bimodal character of gneiss 15, 47, 49  
 Gruinard Bay compared Scourie 59, 59, 61
- Blokken gneiss 236, 242, 242
- Brazilian marbles, geochemistry 121
- British Columbia, removal of melts 105
- bronzite picrite dykes *see* picrite dykes
- brown gneiss, Stoer area  
 metamorphic conditions for 95–6  
 mineralogy **94**, 95  
 partial melting conditions 97, 101
- buffering of metamorphic fluids *see* metamorphic fluids
- Burwell domain 253, 254, 255, 257
- calc-alkaline magma, generation of 52
- calc-alkaline volcanics, E. Hebei Province, China 298
- Caledonian Orogeny **28**, 194, 196  
 foreland 196  
 Lewisian imprint 10–11  
 mylonite formation 39  
 remanence in 207  
 thrusts 10
- Canada, dyke swarms, pre-Grenville Age 217
- Canisp Shear Zone  
 Badcallian deformation 168  
 Inverian deformation 88, 130, 168–9, 168  
 Laxfordian deformation 130, 169–71, 169, 170, 171  
 location 14, 17–18, 129, 130, 165  
 movement of 88, 131–2, 133, 137, 171–2, 172  
 previous work 165–6  
 rock types and structural setting 166–7, 166, 167
- Caozhuang supracrustal association 298, 298, **298**, 299, 299, 300
- carbon dioxide  
 marbles, cause of origin 111  
 metamorphism, heat source 17, 74  
 partial melting, effect on 86, 99–100  
 rare earth elements, removal of 16, 51, 53, 105
- Carnmore antiform  
 Inverian deformation 143  
 Laxfordian deformation 146, 148, 148–9, 149  
 location of 140, 141, 142
- Carnmore shear zone  
 location of 129, 130  
 movement of 131
- cataclasite, Outer Hebrides fault zone **40**, 40, 176, 177, 180, 181  
 microstructure 178, 179  
*see also* mylonite
- Charcot Fjelde dykes, E. Greenland 236, 242
- Charnockites  
 S. Harris, early work 7  
 E. Hebei Province, China 298, **298**, 299, 299, 300, 301, 303  
 E. Greenland 236, 242  
 Madras 48, 49, 51, 105
- Churchill foreland zone, Labrador 253–4, 253, 256
- Churchill Province, Labrador  
 Archaean boundary 251–2, 252, 253–5, 253, 256  
 compared with Greenland Archaean boundary 255–7, **256**, 257
- Claisfean shear zone 17, 18
- Cleitichean Beag dykes 35
- Clospet granite, production of 51
- CMASH model system, use of 97–100, 98, 99
- Cnoc an t'Sidhean supracrustals 93, **94**
- COCORP profiles, Appalachians, compared 193, 202
- Coll/Tiree marbles  
 geochemistry 117  
 limestone-dolostone interbanding 115  
 lithology and mineralogy **112**, 113
- Columbia River basalt province, geochemistry, compared Scourie dyke 231
- cooling  
 isobaric—Napier complex, Antarctica 22, **287**, **289**, 290–1, 292, 293–4  
 modelling 76, 77–8, 77, 77  
 rates of 82, 88–9, 89
- Corodale gneiss 29, 30, 39, 175, 176, 181
- crush melange, Outer Hebrides fault zone 176, 176, 177, 180, 181
- crustal extension, Scourie dyke emplacement 144
- crustal level  
 Laxfordian deformation, change of 19, 140–1  
 Scourie area compared Gruinard Bay 59–60
- crustal reflective character 199, 200, 201–2
- Dalradian marbles, compared Lewisian 117, 121, **122**
- Darling fault 277, 282
- Deccan basalt province, geochemistry compared with Scourie dyke 231
- density of Lewisian rocks 185, **186**, 187
- dextral strike-slip regime, Laxfordian 20
- dextral transpressional tectonic regime 18, 20, 23, 143, 149, 149, 150
- dextral transtensional tectonic regime 20, 143, 144, 146, 149, 150
- Diabaig inlier  
 dykes 162  
 foliation/lineation distribution 157, 157
- Diabaig shear zone  
 Inverian deformation, evidence of 143, 143  
 Laxfordian deformation, evidence of 131, 145, 146  
 location of 129, 130, 140, 141, 142
- diorite dykes, Outer Hebrides 34
- dioritic gneiss  
 central region 14  
 Greenland 236, 261  
*see also* tonalite
- dip-slip movement, Canisp Shear Zone 169, 171–2
- dolerite dykes  
 mineralogy 224, **225**, **226**, **230**  
 occurrence of 218  
 other Scourie dykes compared 229–30, 229, 230  
 petrogenesis 230–1, 232  
 trace element geochemistry 226–7, 226, 227  
 transverse variations and margins 224, 227, 228  
 Antarctica 22, 231, 285, 292

- E. Greenland 235, 236–7, 236, 237, 239  
 compared Scourie dykes 235, 244–5  
 Labrador 253, 254  
 compared Greenland 255, **256**, 257, 257  
 W. Australia 281, 282
- dolomitization  
 effect of Mg/Ca ratio 117  
 of limestones 117–19
- dolostones  
 geochemistry 116, 116, 118  
 limestone/dolostone relationship **114**, **115**, 117–19  
 relationship to other marble groups 121, **121**, **122**
- DRUM seismic profile 193, 194  
 Moho and upper mantle, evidence of 199–200  
 Outer Hebrides fault zone, evidence of 199
- Drumbeg area  
 garnet-pyroxene thermometry 84  
 pyroxene thermometry 83, **84**
- ductility, of crust  
 by underplating 16, 49  
 horizontal tectonics 54
- Dugel gneiss 278, 279
- dykes  
 discussion of 217  
*see also* dolerite dykes, norite dykes, olivine gabbro dykes, picrite dykes, Scourie dykes, 'Younger Basic' suite
- Eastern Hebei Province, China  
 age of 299  
 banded iron formation 298, 298, 299, 299, 300, **300**, 301, 302, 303, 304  
 metamorphic conditions 48, 301–3  
 rock types and petrography 297, 298, **298**, 299, **299**, 301  
 structure 298, 300–1
- electrical conductivity  
 Lewisian rocks **186**, 190, **190**  
 NW Britain, values compared 188, 190, 191  
 porosity, effect of 185
- Enderby Land *see* Napier complex
- epidiorite dykes 18
- Eu anomaly  
 amphibolites, W. Greenland 266, 267  
 granulites 16, 47, 48, 48, 49, 49  
 explained by granite production 50  
 Gruinard Bay 63, 65, 65, 66, 66, 67, 67, 68, 68, 69, 69
- feldspar thermometry 87–8, 87
- Fiskenaesset complex 261, **261**, 262, 262  
 petrogenesis 265, 272–3, 273  
 source magma 270, 273
- Flaggy gneisses, Inverian  
 Torridon shear zone 153, 154
- Flannan complex 195
- Flannan Thrust, seismic reflection 198, 199, 200
- Flannan Trough, formation of 196–7
- fluids *see* metamorphic fluids
- Foindle zone 17
- folding  
 mainland 20, 21, 22, 81, 88  
 Outer Hebrides 36, 37, 43  
 E. Hebei Province, China 300
- E. Greenland 236, 237  
 Napier complex, Antarctica 286–7, 288, 289, 291  
 foliation 23, 34, 37  
 Canisp Shear Zone 166–7, 168  
 W. Greenland 23
- gabbro  
 S. Harris igneous complex 19  
 E. Hebei Province 299, 301
- gabbro-anorthosite complex  
 W. Australia 278, 281  
 W. Greenland *see* Fiskenaesset complex
- gabbro-ultramafic complex  
 garnet-pyroxene thermometry 83, 84  
 Gruinard Bay 61, 81  
 mineral assemblage 82–3  
 Scourie area 58, 59, 61, 81  
 two-pyroxene thermometry 83–4, **84**
- Gairloch metasediments  
 age of 111  
 limestone/dolostone interbanding 115  
 mineralogy 113, **112**, 121  
 skarns **120**
- Gairloch shear zone  
 Inverian deformation 143, 143  
 Laxfordian D<sub>1</sub> deformation 131, 144, 145, 146  
 Laxfordian D<sub>3</sub> and D<sub>4</sub> deformations 141, 148, 149, 149  
 location of 129, 130, 140, 141, 142  
 model of 132  
 relationship to Langavat shear zone 135, 136  
 garnet-pyroxene thermometry and barometry 82, 83–4, 84, **85**, 88, 89
- Gascoyne mobile belt 277, 278  
 geophysical surveys *see* seismic surveys
- Glen Dessarry marble 109, 110
- Glen Urquhart marble 109, 110
- Glenelg marbles  
 age of 111  
 lithology and mineralogy **112**, 113
- granite intrusions  
 Outer Hebrides 20, 28, **28**, **30**, 33, 34, 37, 38  
 Torridon shear zone 154  
 W. Greenland 8
- granitic gneiss  
 Gruinard Bay 57, 59  
 geochemistry **62**, **63**, 66, 67  
 petrogenesis 69, 69, 70  
 petrography 60–1  
 rare earth distribution 67, 68, 69  
 trondhjemites, relationship to 66, 67  
 metamorphism 81, 85–7, 87–8, 87  
 Scourie area 81  
 E. Hebei Province, China 298, **298**, 300, 301  
 Greenland 236, 245, 261  
 Labrador 254  
 W. Australia 277, 278, 280, 281
- granodioritic gneiss  
 Gruinard Bay 67  
 Outer Hebrides 31, **32**  
 E. Greenland 236  
 E. Hebei Province, China 300, 301
- granulite-facies metamorphism  
 E. Hebei Province, China 48, 301, 302, 303

- granulite-facies metamorphism (*cont.*)  
 Greenland 236, 237, 240, 242, 248, 250  
 Labrador 253, 254  
   compared Greenland 256  
 Napier complex, Antarctica 22, 285, 287–90, **287**,  
**289**, 291–2, 292, 293  
 W. Australia 277, 280–1, 281, 282, 283  
   *see also* Badcallian granulite-facies metamorphism  
 graphite schists, W. Greenland 238  
 gravity surveys 185, 187–8, 189  
 gravity variations, Outer Hebrides 29, 31  
 Greenland *see* Ameralik dykes, Amitsoq gneisses,  
 Malene metavolcanic amphibolites,  
 Nagssugtoqidian mobile belt  
 Greenland marbles, geochemistry of 121  
 greenschist facies metamorphism  
 mainland, Laxfordian 22, 128, 140, 148, 207  
 Outer Hebrides fault zone 42  
 E. Greenland 242  
 Labrador 256  
   compared W. Greenland **256**  
 W. Australia 281–2, 283  
 greenstone belts 8, 11  
 E. Hebei Province, China 303  
 W. Australia 277, 278, 279, 280, 280, 281  
 Grenville deformation 22, 128, 149  
 Outer Hebrides fault zone 197–8  
 grey gneiss 8  
 Assynt region 166, 166–7  
 E. Hebei Province, China 299, 299, 300, 301  
 Gruinard Bay  
 geochemistry 16, 50, 61–3, **62**, **63**, 64, 65–7, 65, 66,  
 67, 68  
 geological setting 57–9, 58  
 metamorphic conditions 81–2, 83, **84**  
 petrogenesis 63–4, 68–9, 69, 69–70  
 petrography 14, 60–1, 81, 82  
 Scourie area, compared 59–60, 59  
 heat producing elements *see* lithophile elements  
 Hebei Province, China *see* E. Hebei Province  
 hematite  
 crystalline anisotropy 213  
 remanence record 205, 207, 211  
 hornblende fractionation, effect on rare earth  
 distribution 48, 52, 65–6, 67  
 hornblende pods, tonalites, Gruinard Bay 58, 65, 66  
 Hudsonian Orogeny, Labrador 251, 252  
 crustal shortening caused by 256  
 Churchill foreland zone, effect on 253  
 Komaktorvik zone, effect on 254, 255  
 Nain Province, effect on 253  
 hydrous fluids *see* metamorphic fluids  
 hypersolvus granites, composition of 87, 87  
 ilmenite-magnetite intergrowths, iron-titanium  
 thermometry and bathometry 85–7  
 Indian Archaean terrain  
 petrogenesis 48–9  
 Interior Churchill orogen, Labrador 253, 254–5, 255,  
 256  
 Inverian  
 deformation-shear zones **15**, 17–18, 93, 128, **128**,  
 129, 130, 131  
 Canisp 17–18, 88, 129, 165, 168–9, 168, 169,  
 171, 172  
 kinematic model 142–4, 143, 144  
 Torridon 155, 159  
 Nagssugtoqidian, Greenland, compared 23, 244  
 significance of 10  
*see also* amphibolite facies metamorphism  
 iron-titanium oxide thermometry and barometry 85–  
 7, 89  
 Isua supracrustal association, W. Greenland 236  
 Itivdleq shear zone 249–50, 250, 250–1, 251  
 Jequie complex, Brazil, lithophile depletion 49, 51  
 Kabbaldurga Quarry, S. India, evidence of CO<sub>2</sub> fluid  
 element removal 51  
 Kangâmiut dykes, W. Greenland  
 Labrador, compared 255, **256**  
 Nagssugtoqidian mobile belt 249, 249, 250, 250,  
 251, 251  
 Kangerdluk-Ugpiik suite, Malene metavolcanics, W.  
 Greenland 266, 272  
 Karoo basalt province, geochemistry, compared  
 Scourie dykes 231  
 Katarchaeon continental crust, Scourian accretion  
 8–9  
 Kenmore shear zone 153, 154, 162  
 Laxfordian D<sub>1</sub> deformation 131, 144, 146  
 Laxfordian D<sub>3</sub> deformation 141, 148  
 model of 133  
 Kenoran Orogeny, Labrador 251  
 Ketilidian, Greenland 9, 10, 23  
 Komaktorvik zone 253, 254, 255, 256  
 Komatiitic basalt types, Malene metavolcanics, W.  
 Greenland **263**, 265, 266–9, 266, 267, 272  
 Kylesku Group, density 185, 187  
 lacunae–low strain areas, Torridon shear zone 154,  
 155, 157–8, 158–9, 161  
 Lake Harbour Group 252, 255, 256  
 laminar flow conditions, dyke intrusion 228, 232  
 Langavat belt 29, 33  
 Langavat shear zone 129, 130, 134  
 movement 132–5, 134  
 relationship to mainland 135–6, 135  
 Laxford shear zone  
 description of 88, 129–30, 131  
 location 129, 130  
 movement 129–30, 132, 133  
 relationship to Langavat shear zone 135, 136  
 Laxfordian 10, **15**, 19–22, 21  
 Gruinard Bay 57  
 kinematic model 140–1, **141**  
 D<sub>1</sub> deformation 144–6, 146, 147  
 D<sub>2</sub> deformation 146–8, 146, 147  
 D<sub>3</sub> deformation 148–9, 149  
 D<sub>4</sub> deformation 149, 149  
 Outer Hebrides **28**, 36–43, **36**  
 shear zones 23, 93, 129, 131–2, 150  
 Canisp 165, 168, 169, 169–71, 170, 171, 171–2,  
 172  
 Laxford 129–30  
 Outer Hebrides 132–5  
 Torridon 155, 159, 160, 161–2, 161, 162

- Laxfordian (*cont.*)  
 Nagssugtoqidian, Greenland, compared 23, 244, 244
- Leotheid dyke, transverse modal variations and margins 219, 220–1, 221
- Letterewe gneiss, emplacement of 148, 148
- Letterewe synform  
 deformation of 146, 148  
 location of 141, 142
- leucogneiss  
 Stoer area  
 mineralogy 93, 94, 102  
 petrogenesis 101, 103, 103  
 E. Greenland 242, 243
- leucogranite, Outer Hebrides 37
- Leverburgh belt 29, 33
- Lewisian  
 early work 7–8  
 location 14  
 post-Laxfordian, effect on 10–11  
 relationship to Precambrian Europe and America 9–10  
 Scourian accretion 7, 8–9, 11
- lherzolite garnet mantle source 269–71, 270, 272
- limestone, chemical division 113  
*see also* metasedimentary marbles
- lineations, elongation, guide to shear movement 127, 129, 130, 131
- lithophile elements  
 continental flood basalts 231  
 explained by  
 flushing of CO<sub>2</sub> rich fluids 16, 51–2, 53  
 melt extraction 16, 50–1, 53  
 granulite facies metamorphism, removal in 16, 22–3, 49–50  
 amphibolite facies, compared 47, 50, 50  
 Scourie dykes 18, 227, 227, 230–1
- lithosphere thickening *see* tectonic thickening
- lithospheric thinning *see* tectonic thinning
- Loch Maree fault, location of 140, 141
- Loch Maree Group  
 emplacement of 19, 21, 128, 128, 141, 143, 144, 144, 148, 148  
 location 14  
 Greenland and Labrador equivalents 23
- Loch Maree Group marbles  
 age of 111  
 compared other Lewisian marbles 118, 119, 121, 122, 123, 123, 124  
 location 110  
 mineralogy and lithology 112, 113, 116
- Loch Skerrols Thrust 194, 197
- Lochalsh fold 159–60
- Lochinver monoform 165, 166, 166, 167, 168, 168, 169, 170, 171
- lower detachment shear zone 132, 132, 133  
 Outer Hebrides 135
- mafic-ultramafic complexes 13, 14, 45, 46  
 petrogenesis 48–9  
 rare earth abundance 47, 48  
 Antarctica 286  
*see also* ultramafic gneiss
- maghemite  
 remanence in 206
- shape anisotropy 213
- magnetite-talc rock 110, 120
- magnetism  
 factors influencing 185, 187, 205  
 surveys 186, 187–8, 189  
 W. Australia 282  
*see also* palaeomagnetic record
- magnetite  
 remanence in 185, 187, 205, 206, 207  
 shape anisotropy 213
- Malene metavolcanic amphibolites, W. Greenland 261, 261, 262, 262  
 Ameralik dykes, compared 264, 265, 266, 268, 271–2, 271, 272  
 Fiskenaeset metagabbro-anorthosites, compared 273, 273  
 geochemistry  
 Ivisártoq 262–5, 262, 263, 264, 265, 267, 268  
 Ravns Storø 262, 265–6  
 petrogenesis 266–71, 269, 270
- Malene supracrustals, W. Greenland 247, 248  
 Lewisian marbles, compared 111
- Manfred complex, W. Australia 278
- marbles  
 age of 111  
 ancient carbonates, other, compared with 117, 121, 122  
 chemical subdivision 113  
 distribution of 29, 109, 110  
 geochemistry 115–17, 116, 118, 119, 120  
 limestone/dolostone relationship 114, 115, 117–19  
 lithology and mineralogy 112, 114, 115, 115  
 origin, evidence for 110–11  
 tectono-metamorphic 110, 110, 120  
 E. Greenland 238  
 E. Hebei Province, China 298, 304  
 Labrador 255
- Meeberrie gneiss, W. Australia 278, 279, 280–1  
 melt migration  
 energy transfer for metamorphism 75
- mesoperthite feldspars, evidence of  
 'dry' conditions 287  
 high temperatures 87, 87, 88
- metabasic rocks 8  
 age 33  
 metasedimentary rocks, association with 33  
 mineralogy 29, 31  
 Outer Hebrides 28, 29, 30  
 Greenland, compared 33
- metadolerite dykes  
 Canisp Shear Zone 169  
 Outer Hebrides 35
- metagabbros, S. Harris, magnetic anomaly 187
- metamorphic fluids  
 lithophiles, enrichment of 31  
 partial melting, effect on 97–100, 99, 101, 103  
 retrogression, effect on 86–7, 181, 231, 253
- metapelites  
 significance of 294  
 E. Hebei Province, China 298, 304
- metasedimentary gneiss 8, 13, 16  
 Caledonian foreland 196  
 mafic-ultramafic bodies, association with 15, 46, 47

- metasedimentary gneiss (*cont.*)  
 mainland 81  
   metamorphic conditions 83–4, 84  
   melting, evidence of 50–1  
 Outer Hebrides 28, 29, 30, 33  
   metabasic rocks, association with 33  
   Greenland, compared 33  
 Antarctica 286, 288  
 E. Greenland 235, 237  
 E. Hebei Province, China 298, 299, 300, **300**, 301  
 Labrador 253, 255  
 W. Australia 277, 279
- metasomatism  
 due to CO<sub>2</sub> movement 51  
 E. Hebei Province, China 299
- microdiorite dykes, Outer Hebrides **31**, 33–4, 38–9
- migmatization 8, 20  
 E. Greenland 244  
 Outer Hebrides 29, 33–4  
 Torridon shear zone 153–4  
 Labrador 253  
 Napier complex, Antarctica 294
- Minch Basin 196–7  
 formation of 195, 197–8
- Minch Fault  
 generation of 39, 43  
 movement of 135, 135
- Moho  
 depth variation 202  
 reflective properties of 199–200, 200
- Moine Outcrop, Lewisian marbles 109, 110
- Moine schists, Lewisian inliers in 196
- Moine Thrust zone 181, 194, 196, 201, 201, 202
- MOIST seismic profile 193, 194–6, 194, 197
- Moho and upper mantle, evidence of 199–200  
 moine thrust, evidence of 201, 201
- monzodiorite dykes, Outer Hebrides 34
- monzonite  
 E. Hebei Province, China 299, 301  
 W. Australia 278
- mylonite zone  
 Canisp Shear Zone 169–70  
 characteristics of 180–1  
 Outer Hebrides fault zone 39–41, **40**, 42, 43, 181  
 Labrador 254, 255  
 W. Australia 277  
*see also* phyllonites
- myrmekite, E. Hebei Province, China 299, 301
- Nabberu Basin 277, 278
- Nachvak granulites 253
- Nagssugtoqidian, Greenland 8, 10  
 Hudsonian, compared 23, **256**  
 Inverian and Laxfordian, compared 23, 244  
 Nag. 1 deformation 236–7, 240, 241, 243, 249–50, 250–1, 251  
 Nag. 2 deformation 242, 242, 243, 250, 251, 251
- Nagssugtoqidian mobile belt  
 E. Greenland  
   Archaean gneiss complex 236, 236, 237, 238  
   Blokken gneiss 240–2, 242  
   Nag. 1 deformation 236–7, 239, 240, 241  
   Nag. 2 deformation 242, 242, 243  
   post-tectonic intrusions 242  
   supracrustal rocks 238–40  
 W. Greenland  
   boundary Archaean gneiss 247–51, 248, 249, 250, 251  
 Labrador boundary compared 255–6, **256**, 257  
 Lewisian  
   compared 23, 235, 235, 244–5, 244  
   relationship to 132, 136, 137
- Nagssugtoqidian supracrustals, Greenland,  
 correlation with Lewisian marbles 111
- Nain Province, Labrador 248, 251, 252–3, 252, 253, 255, 256
- Napier complex, Antarctica  
 cooling **287**, **289**, 290–1, 292, 293–4  
 deformation 286–7, 291  
 dykes, compared Scourie dykes 231  
 early work 285  
 lithology 285–6  
 location 285  
 metamorphic conditions 22, 285, 287–90, **287**, **289**, 291–2, 292, 293  
 pre-deformation history 292–3
- Ness assemblage, Outer Hebrides 29
- Ness shear zone, location 129, 130
- Nordre Stromfjord shear zone 157
- Noritic dykes  
 mainland  
   compared other Scourie dykes 229–30, 229, 230  
   mineralogy 224, **225**, **226**, **230**  
   occurrence 224  
   petrogenesis 18, 230–1, 232  
   trace element geochemistry 226–7, 226, 227  
   transverse variations and margins 224, 227, 228
- Outer Hebrides 35
- normal movement, Outer Hebrides fault zone 176, 181–2
- Nük gneiss, W. Greenland 217, 248, 261, 272  
 Labrador, equivalent in 253
- Oaxacan granulites, Mexico, feldspar composition 87
- olivine  
 Malene metavolcanics, W. Greenland, importance in evolution 268–71, 269, 270, 272  
 Scourie dykes, variability of—  
   margins 228  
   olivine gabbro dykes 222, **222**, 223, 223  
   picrite dykes 218, 220, 220, 221, **221**
- olivine gabbro dykes  
 compared other Scourie dykes 229–30, 229, 230  
 mineralogy 222–3, **225**, **226**, **230**  
 occurrence 221–2  
 petrogenesis 18, 230–1, 232  
 trace element patterns 226–7, 226, 227  
 transverse variations and margins **222**, 223–4, 223, **224**, **225**, 227, 228, 228–9, 232
- orthogneiss  
 Stoer area 93, **94**, 96, **96**  
 Antarctica 286, 287, 288  
   precursors of 292–3, 294  
 E. Hebei Province, China **300**, 301, 302, 303  
 Labrador 253
- osumilite, evidence of 'dry' conditions 287, 288

- Outer Hebrides  
 chronology **28**  
 Laxfordian 20–2, 21, 35–9, **36**, 38  
 lithology 27–33, **28**, 29, **31**, **36**  
   E. Greenland, compared 244  
 metamorphic conditions 29, 33, 35  
 palaeomagnetism 209–11, 210, 211, 212, 213  
 Scourian intrusives 19, 34–5  
*see also* Langavat shear zone, South Harris igneous complex
- Outer Hebrides fault zone **22**  
 age 196, 197–8, 202  
 base of 198–9  
 location 39, 175, 175, 176  
 mainland shear zones, relationship to 46  
 metamorphic conditions 39–41, 43, 175, 175, 176, 181, 187  
 microstructure 177–9, 178, 180  
 mylonite zone 180–1  
 pre-existing boundary 187  
 rock types 39–41, **40**, 42, 43, 41, 176–7, 176  
 thrust or fault zone 175–6, 181–2, 197, 198, 199
- Outer Hebrides Thrust *see* Outer Hebrides fault zone
- Outer Isles Thrust *see* Outer Hebrides fault zone
- palaeomagnetic record  
 age of 205–6, 211–12  
 mainland  
   central and north 206–7, 207  
   south 207–9, 208, 209  
 Outer Hebrides 209–11, 210, 211, 212, 213  
 shape anisotropy, effect of 212–14
- paragneiss *see* metasedimentary gneiss, marbles
- partial melting  
 batch melting 103–4, 104  
 CMASH model system 97–9, 98, 99  
 conditions for 70, 88, 96–7, 99–100  
 hydration state, importance of 102–3  
 melting in—  
   amphibolites 68–9, 69, 69–70  
   metabasites 97–100  
   supracrustal lithologies 101  
   tonalitic gneisses 100–1  
 lithophile distribution, effect on 50, 53, 54, 105  
 thermal models for metamorphism 75, 76  
 trace element distribution, effect on 104–5
- pegmatites  
 post-Laxfordian 10, 206  
 Scourie and Laxfordian 8, 34–5, 46, 165, 244  
 E. Greenland 237  
 E. Hebei Province, China 299, 301, 303
- pelitic schists, E. Greenland 238
- peridotites, Scourie compared Gruinard Bay 60
- permeability  
 Lewisian **186**, 190, **190**  
 porosity, effect of 185, 187
- Perth Basin 277
- phyllonites  
 Outer Hebrides fault zone 39, **40**, 41, 176, 176, 177, 180, 181  
   microstructure 41, 179, 180  
 W. Australia 282
- picrite dykes  
 compared other Scourie dykes 229–30, 229, 230  
 late stage products 221  
 mineralogy 218–19, **225**, **226**, **230**
- Outer Hebrides 35  
 petrogenesis 18, 230–1, 232  
 trace element patterns 226–7, 226, 227  
 transverse variations and margins 219, **219**, 221, **225**, 227, 228, 232  
   Beannard dyke 219–20, 220, 228  
   Leothaid dyke 220–21, **221**
- picritic lava 231
- piedmont fan, Torridon 197
- Pilbara craton, W. Australia, age of 277, 278
- pinch-and-swell structures, dykes 237, 244–5, 250
- plate tectonics 136, 137, 150, 257, 257
- Poisson's ratio  
 NW Britain, values compared 191  
 variation in 188
- porosity 190, **190**  
 effect on physical properties 185, 187  
 NW Britain, values compared 191
- pressure–temperature conditions *see* amphibolite facies, granulite facies, greenschist facies, partial melting
- 'proto-Atlantic' 196
- psammite, W. Australia 281–2, 282
- pseudotachylite  
 mainland 22, 149  
 Outer Hebrides 39–41, **40**, 42, 175, 179, 181  
 Labrador 254
- pyroxene  
 analysis of Scourie dykes **225**, 229, 229  
 dyke margins 227–8, 232
- pyroxene-granulites  
 magnetization of 187  
 E. Hebei Province, China 298, 301  
 W. Australia 281
- pyrrhotite, remanence in 206
- Q-normative, chemistry metabasics 29, **31**, 34
- Qianan gneisses, E. Hebei Province, China 297, 298, **298**, 302
- Qianxi gneisses, E. Hebei Province, China 297, 298, **298**
- Qôrqut granite, W. Greenland, production of 51
- quartz-dolerite dykes *see* dolerite dykes
- quartz-epidiorite dykes, Canisp Shear Zone 166, 167, 168, 169
- quartz-magnetite, W. Australia 279
- quartz-tholeiitic dykes *see* dolerite dykes
- quartzite  
 Outer Hebrides 29  
 E. Hebei Province, China 298, 301, 304  
 Labrador 255  
 W. Australia 279–80, 281–2, 282
- quartzo-feldspathic gneiss  
 feature of pre-Cambrian gneiss terrain 235, 244  
 Outer Hebrides 28, 29, 31, 32, 33  
 E. Hebei Province, China 300, 301  
 W. Greenland 236, 248  
*see also* granitic gneiss, tonalites, trondhjemites
- radiogenic causes of metamorphism 73, 76, 78
- Raggatt Series, E. Antarctica 286

- Ramah Group supracrustals 252, 253–4, 255  
 Loch Maree Group, compared 23  
 W. Greenland, compared 256, **256**  
 ‘ramp-flat’ geometry, shear zones 129, 131–2, 132, 133, 135, 146–7  
 Torridon shear zone 153, 157, 157, 159, 161, 162  
 rare earth element distribution  
 Gruinard Bay  
 amphibolites 61–4, **62**, 65, 69  
 quartzo-feldspathic gneiss **62**, 65–6, 65, 66, 66, 67, 67, 68, 68, 69, 69  
 marbles 116, 116, **118**, **119**, **120**, 121  
 silicic and mafic gneisses compared  
 Archaean 47–9, 48  
 post-Archaean 49, 49  
 Scourie dykes 18  
 E. Hebei Province, China 302  
 W. Greenland 266, 267  
 Rayner complex, E. Antarctica  
 location of 285, 285  
 underplating of 294  
 remanence  
 acquisition by metamorphic rocks 205  
 age of 205–6  
 fabric, effect of 212–14  
 Laxfordian or post-Laxfordian uplift 211–12  
 mainland  
 central and north 206–7, 207  
 south 207, 207–9, 208, 209  
 Outer Hebrides 209–11, 210, 211, 212  
 pre-Laxfordian, south mainland 207, 209  
 retrogression  
 Inverian 18, 88–9, 89, 90, 95–6, 95, **95**  
 isobaric cooling, Napier Complex, Antarctica compared 294  
 Laxfordian 140, 148  
 Hudsonian, Labrador 253  
 Nag. 1, Greenland 237, 240, 249, 250  
 Rhiconich Group, density 185, 187  
 Rinkian mobile belt, N. Greenland  
 Outer Hebrides, compared 244  
 Nagssugtoqidian mobile belt, Greenland, compared 238, 242  
 Rockall Trough 194, 196  
 Rosemarkie marble 100, 109  
 Ruadh Mheallan belt 131, 140, 141, **141**  
 Badcallian structures 153  
 dykes 162  
 Inverian deformation 143, 143  
 Laxfordian deformation 144–5  
 rutile, retention of elements 53
- Saglek dykes, Labrador 252  
 Sarfartup nuna supracrustals, W. Greenland 256  
 Loch Maree Group, compared 23  
 Scardroy sheet, marble inliers 109, 110, 111, **112**, 113, **114**  
 schistose amphibolite dykes, E. Greenland 237, 242  
 Scourian  
 accretion of 7, 8–9, 11  
 gneiss, origin of 15–16  
 granulites 9, 10  
 Outer Hebrides 28, **28**, 29–36  
 see also Badcallian granulite facies, Inverian, Scourie dykes  
 Scourie area, gneiss compared Gruinard Bay 59–60, 59, 61, 81–2  
 Scourie dykes  
 age of **15**, 46, 88, 217  
 distribution of 18, 218  
 emplacement of 8, 18, 21, 143, 144, 144, 145  
 magma source 18–19, 230–1, 232  
 margins of 227–8, 232  
 metamorphic conditions 8, 76, 89, 89, 95, **95**, 205–6, 227, 231  
 mineral chemistry variation **225**, 229–30, 229, 230  
 mineral distribution, asymmetrical 228–9  
 petrological groups 218  
 strain indicators in gneiss 127  
 time markers 7, 57, 93, 128, 129, 141, 155, 155  
 trace element geochemistry 18, 226–7, 226, 227  
 see also dolerite dykes, noritic dykes, olivine gabbro dykes, picrite dykes  
 seismic surveys 8  
 crust and upper mantle, evidence of 199–200, 200, 202  
 factors influencing 185, **186**, 187  
 geology and structure, evidence of 194–9, 195, 198, 199  
 Lewisian basement, evidence of 201, 201, 202  
 lines of 189, 193–4, 194  
 NW Britain, Lewisian analogues 188–90, 191  
 shear zones  
 deformation chronology 88, 89, 89, 128, **128**, 129  
 distribution of 15, 17–18, 20, 129, 130, 196  
 mainland 127, 129–32, 132  
 lower detachment zone 18, 34, 132, 132, 133  
 Outer Hebrides 34, 35, 36, 132–5, 134  
 relationship to mainland 135–6, 135  
 role of 9  
 Scourie dykes, relationship to 18, 217  
 shear heating, effect on metamorphism 75  
 E. Hebei Province, China 299, 299, 301, 303  
 Greenland 237, 249–50, 250, 250–1, 251  
 Labrador 253, 254  
 compared Greenland 18, 255, **256**, 257, 257  
 N. Atlantic region, model for 23, 136, 137  
 see also Canisp Shear Zone, Outer Hebrides fault zone, Torridon shear zone  
 SHET seismic profile 193–4  
 Shinness marble/gneiss succession 109  
 lithology and mineralogy **112**, 113  
 Shuichang supracrustal association, E. Hebei Province, China 298, 298, **298**, 300, 301, 304  
 Sinian suberathem, E. Hebei Province, China 298, 298  
 skarns 113, 113, **120**  
 nodules 115, **115**  
 South Harris granulites 19, 133, 134  
 South Harris Igneous Complex  
 age of **28**, 211  
 emplacement of 14, 19  
 magnetic anomalies 187  
 metamorphic conditions 33  
 metamorphism of 211  
 palaeomagnetic record 209, 210, 210, 211, 212, 213



- South Harris marbles  
 age of 111  
 lithology and mineralogy **112, 113, 114**  
 texture of 115
- South Harris shear zone *see* Langavat shear zone
- sphene, sink for rare earth elements 53, 67
- Stoer area  
 deformation of 93, **94**  
 element depletion 105  
 location 93  
 partial melting 96–105, **96, 98, 99, 102, 103, 103**  
 petrology 93, **94, 102**  
 metamorphic conditions 93, 94–6, **95, 95**
- strike-slip regime  
 Laxfordian 23, 149, **149, 150, 171–2**  
 Labrador 23
- subsolus granites, composition of 87, 87
- syenogranite, W. Australia 278
- symplectite formation 83, 84, 85, 88, 89–90, 89
- Tasiuyak gneiss 254, 255
- tectonic thickening, effect on metamorphism 17, 75–8, 95, 283, 290, 293
- tectonic thinning, effect on metamorphism 74–5
- tholeiite  
 Malene metavolcanics, W. Greenland **263, 265, 266, 267, 268–9, 272**  
 origin Scourie gneiss 15
- thrusting  
 Moine Thrust zone 181, **194, 196, 201, 201, 202**  
 Outer Hebrides fault zone 175–6, 181, 182  
 N. Atlantic Archaean boundary 8, 23, 256, 257  
 W. Australia 282, 283
- Tollie antiform 20, 131, **140, 141, 142, 146, 148–9, 149**
- tonalitic gneiss  
 distribution of 13, 14, 45, 46  
 geochemistry **62, 63, 65**  
 lithophile depletion 22–3, 52, 53  
 metamorphic conditions 83  
 occurrence  
 Gruinard Bay 57, 58, 59, **59, 81**  
 Outer Hebrides 19, 29, 211  
 Scourie area 81  
 Stoer area 93, **94, 102**  
 Torridon shear zone 153, 154, 155  
 E. Greenland 236, 242  
 E. Hebei Province, China 298, **298, 299, 300, 301, 302, 303**  
 partial melting 100–1, 103, **103, 103**  
 petrogenesis 48, 49, 52, 53, 68, **68, 70**  
 post-Archaean equivalents 49, **49**  
 rare earth distribution 15–16, 47–8, **48, 65–6, 65**
- Torridon antiform  
 Laxfordian deformation 148–9, **149**  
 location of **140, 141, 142**
- Torridon shear zone  
 evolution 161–2  
 geometry 157–9, **157, 158**  
 explanation of 159–60, **159, 160, 161**  
 structure 153–5, **154, 155, 156**  
 lithology 153–4
- Scourie dykes 154, 155, **155, 156, 161, 162–3**
- Torridonian sandstone, effect on shear plane 160, 161
- Torridonian sequence, NW Scotland 90, 196
- trace element patterns  
 continental flood basalts 231  
 partial melting, effect of 104–5  
 Scourie dykes 226–7, 226, 227, 230–1, 232
- transpressional regime  
 mainland  
 Inverian 131, 137  
 late Laxfordian 132, 137  
 Outer Hebrides, Laxfordian 133
- transtensional regime, early Laxfordian 131, 137
- trondhjemitic gneiss 13, 14, 57, 58–9, 58  
 geochemistry **62, 63, 66, 66, 67**  
 metamorphic conditions 17, 85–7, 88  
 petrogenesis 68, 69, 70  
 petrography 60  
 rare earth distribution 15–16, 66–7, **66, 67**
- Scourie, compared Gruinard Bay 59–60, 81, 82
- Torridon shear zone 154
- veins  
 amphibolites, association with 67  
 partial melting origin 97–100, 98, 99, **103, 103**
- turbulent flow conditions, dyke intrusion 228, 232
- Tula Series, Antarctica 286
- Uivak gneiss, Labrador 252, 253
- ultracataclasite  
 microstructure of 177–9, **178**  
 Outer Hebrides 175, **176, 177, 181**
- ultramafic dykes  
 Canisp Shear Zone 166, 167, **168, 169**  
 Outer Hebrides 34
- ultramafic gneiss, Gruinard Bay 57, 58, 59  
 geochemistry 61, 63, **64**  
 petrography 60
- ultramafic pods, Assynt region 166
- ultramylonite, Outer Hebrides 39, 41, 42
- underplating, tonalitic magma 16, 49, 54
- Upemavik supracrustals, Labrador 252–3, 254, 255
- uplift 39, 43, 90  
 rate of 9, 10, 76  
 Antarctica 293
- Upper detachment shear zone 132, **132, 133**
- Western Shield, Australia 277, 278
- Western gneiss Terrain, Australia 277, 278, 280, 283  
 N. sector 278–81  
 S. sector 281
- white mica layer, partial melting of 101
- WINCH seismic profile  
 crustal reflection 199, **200**  
 line of 189, 193, **194**  
 Moho reflection 199–200  
 Outer Hebrides fault zone 197, 198, **198, 199**
- Yilgarn Block, W. Australia, age of 277, 278, 283  
*see also* Western gneiss Terrain
- 'Younger Basic' (Scourie dyke) suite 28, **31, 35–6, 37**