

Index

- Acraman impact ejecta layer 53–4, 117, 123, 126–9, 130–2, 425–7
- acritarchs
- Amadeus and Officer Basins 119
 - biostratigraphy 115–25, 130–2
 - Australian correlations 130–2
 - composite zonation scheme 119, 131, 132
 - India 318–20
 - Ireland 289
 - Spain 232
 - sphaeromorphid 324
- Adelaide, Hallett Cove 68
- Adelaide Rift Complex 115–22, 425
- correlations with Officer Basin 127
 - dating (Sr–Rb) 140
 - generalized time–space diagram, correlations
 - between tectonic units 120
 - location maps 116, 118
 - SE sector cumulative strata thickness 139
 - stratigraphic correlation with Officer Basin 127
 - Stuart Shelf drill holes, correlations 117
 - Sturtian (Umberatana) Group 116, 138
 - Umberatana Group 116, 138
- Africa
- Vendian climatic indicators 17
 - see also* Namibia
- Aldanellidae 418
- algal metaphyta, White Sea Region 271–4
- algal microfossils, White Sea Region 271
- Altenfeld Formation, backarc spreading 44–5, 47–8
- Amadeus Basin
- acritarch zonation 119
 - carbon isotope trends 128
 - generalized time–space diagram, correlations
 - between tectonic units 120
 - location map 118
 - succession 124–5
- Anabarella 230
- Anabarites *tricarinatus* 193
- Anabarites–Protohertzina Zone, Precambrian–Cambrian
- boundary, China 145–7, 423–4
- Andean Basin 1–14
- Andiva 158
- Animalia, origins 447–8
- Appendisphaera barbata*/*Alicesphaeridium medusoidum*/*Gyalosphaeridium pulchrum*
- Assemblage Zone 119, 131, 132
- Arabia, Vendian climatic indicators 17
- Archaeobranchia 418–19
- Archaeotrichion lacerum* 430–1
- Archyfasma* 271–4
- Argentina (NW), Puncovicana Formation 1–14
- location map 2
 - stratigraphy 3
- Armorica 36
- armour, and infaunal shelters,
- origins 405–14
- arthropod-type trace fossils, Spain 226–7
- Arumberia banksi* 285, 290
- Aspidella*
- controversy 300, 301–3, 305
 - ecology 303
 - synonyms 302
- Australia
- Acraman impact ejecta layer 53–4, 117, 123, 126–9, 130–2, 425–7
 - carbon isotope chemostratigraphy 126–9
 - correlations of Ediacaran System and Period 18, 115–35
 - Marinoan glaciation 53–4, 126
- Australia, Ediacaran System and Period
- Bunyeroo–Wonoka Formation transition 137–9, 426
 - Centralian Superbasin 118, 125
 - composite zonation scheme 131
 - correlation methods and results 125–32
 - time–space diagram 120
 - Vendian climatic indicators 17
 - See also* Adelaide Rift Complex; Flinders Ranges
- Avalonian assemblages, Newfoundland 237–57, 303–7, 427
- backarc spreading, Altenfeld Formation 44–5, 47–8
- Baliana–Krol Group, NW Himalaya 319
- Barut Formation, Iran 434
- Bayesian analysis
- eumetazoans 357–9
 - Porifera 357–9
- Belarus, Vendian cyclic structure 21
- Beltanella* 324
- Beltanelliformis*
- Australia 427
 - Russia and Ukraine, comparative taphonomy 259–67
 - Spain 224, 228
- Beltanelloides* 4, 8, 271–4
- Bilateria 369–75
- first appearance 126, 157, 447–8
 - gliding symmetry 399
 - small-bodied crown phyla 371–2
- biosiliceous sedimentation, China, earliest
- Cambrian 146, 423–4
- Birmanian basin, Rajasthan 439–41
- Blaini–Krol–Tal Basin, Lesser Himalaya
- carbon isotope chemostratigraphy 85–91
 - geological map 79
 - global correlations 96
 - trace elements 90
- Blainiella* 318
- Blumenau Shear Zone 37
- Bohemian Massif, Saxo-Thuringian Zone (Cadomian basement) 35–51
- geological map 36–8
- Bonney Sandstone, Flinders Ranges 122, 140, 197, 426
- Brachina Formation 426
- Trezona Bore 65–7, 69–70
- Brachina Gorge, Ediacaran GSSP 196
- Bradgatia* 216

- Breakfast Time Creek Member, Rawnsley Quartzite, Flinders Ranges 197–8
- Bunyerroo Gorge 116
- Bunyerroo–Wonoka Formation transition 121–2, 137–9, 426
- Wearing Dolomite 117, 121–2, 137–9
- Buxa Dolomite, Lesser Himalaya
- carbon isotope chemostratigraphy 91–5
- microfossils 97–8
- Ranjit Window 81, 94
- stromatolite assemblages 93–5
- Cadomian orogeny, Saxo-Thuringian Zone 35–51
- backarc basin 41–4
- Cadomian Unconformity 41
- retroarc basin 45, 47–9
- Cadomian unconformity 35–6, 41
- Cadomian–Avalonian active margin, Peri-Gondwana 46
- calcareous tubular fossils 410
- calcsponges, parapyly 355–67
- calcite–dolomite cycles, Namibia 103–13
- Cambrian ‘problematica’ 360–2
- cap carbonates
- calcite–dolomite cycles, Namibia 103–13
- carbon isotope chemostratigraphy 126–9
- rhythmic sequences 27–34
- ‘Snowball Earth’ hypothesis 23
- see also* carbonates
- cap dolostone
- magnetics 69–72
- Nuccaleena Formation 54, 59–61, 67–72, 120–1
- carbon isotope chemostratigraphy
- Australia 126–9
- Birmania basin, Rajasthan 439–41
- delta $^{13}\text{C}_{\text{carb}}$ values, beginning and end of Ediacaran System and Period 427
- Deoban Group 82, 88
- Lesser Himalaya
- Blaini–Krol–Tal sequence 85–91
- Buxa Dolomite 91–5
- Lameri Limestone, Garhwal 86
- Nemakit–Daldynian 440
- and Precambrian oxygenation 362–3
- Proterozoic ocean 362
- carbonaceous compressions 228–30
- carbonaceous microfossils, India 324
- carbonaceous microfossils, White Sea Region 269–75
- carbonates
- delta $^{13}\text{C}_{\text{carb}}$ values 27
- Proterozoic ocean 362
- see also* cap carbonates
- Centralian Superbasin
- Australia 118, 125
- location map 118
- Ceratosphaeridium mirabile*/*Distosphaera australica*/*Apodastoides verobturatus*
- Assemblage Zone 119, 131, 132
- Charnia* fronds
- affinities 251
- Australia 210–12
- Avalon peninsula, Newfoundland 237–57
- ecology/life 248–51
- landmark morphometric analysis 240–8
- stratigraphic setting 238
- taphonomy 242
- traditional interpretation of morphology 239
- quiltlike 400
- Charniodiscus* 158, 199, 211–12, 297–8
- affinities 251
- feeding 377
- holdfast 322
- quilting patterns 388
- Chase Quartzite Member, Rawnsley Quartzite 197
- China
- Doushantuo Formation 151–6
- earliest Cambrian 423–4
- Precambrian, ciliated protozoans 151–6
- Precambrian–Cambrian boundary, radiolarian skeletons 143–9, 424
- Vendian climatic indicators 16–17
- Yunnan, Meishucun section 411
- chirodropids, cubozoans, *Kimberella* 157–79
- Chondrophorina 217–18
- Chondroplon* 214, 217
- Chuar*, Iran 435, 436
- Chuar*–*Tawuia* association 324, 328
- Chuosi Formation, cap carbonates, Namibia 27–34, 103–13
- ciliated protozoans, Precambrian, China 151–6
- Ciliophora
- Precambrian, S China 151–6
- systematic palaeontology 155–6
- climatic indicators
- palaeoclimatic and biotic changes 88
- Vendian Period 15–26
- Cloudina*-bearing sediments, Spain 223–4, 230–2
- Cnidaria 218, 355
- Cochlichnus anguineus* 5, 7
- Conomedusites* 320–2
- Coreospiridae 417
- Corynetis* priapulid 408
- crown phyla, small-bodied bilateria 371–2
- ctenophores 355
- cubozoans 157–79
- see also* *Kimberella*
- cyanobacterial films/mats 197, 269–71
- Deoban Chert 78
- Cyclomedusa* 300, 301, 302, 321
- demosponges, parapyly 355–67
- Deoban Group, carbon isotope chemostratigraphy 82, 88
- Deoban–Gangolihat carbonate belt 78, 88, 95
- Dickinsonia* 158, 198, 199, 292, 323, 443
- carpet-like recliners 389
- feeding 377
- quilting patterns 387–8
- resting traces 394
- Didymaulichnus* *see* *Mattaia*
- Diplocraterion* 409
- discoidal fossils 297–313
- Aspidella* controversy 301–3
- history of research 297–8
- morphology 299–301
- pseudofossils 301

- taphonomy 298–9
 taxonomy 308–9
- dolomite
 calcite–dolomite cycles 103–13
 Gangolihat Dolomite Formation,
 chemostratigraphy 83
 synsedimentary model 110–12
 rip-up clasts 111–12
- Doushantuo Formation, China 151–6
- Eastern Europe, Vendian climatic indicators 16
- Ediacara Member and fauna/assemblage 198–200,
 195, 425–8
 new approach (Morphospace Analysis) 378–9
 preservation 199–204
 Rawnsley Quartzite (Pound Subgroup),
 Flinders Ranges 198–200
- Ediacaran GSSP
 Brachina Gorge 196
 defining 73, 115, 425
 Elatina Formation, rock magnetic experiments 71
 Enorama Creek 54, 55–61, 116
- Ediacaran System and Period
 acritarch biostratigraphy 115–25, 130–2
 composite zonation scheme 131, 132
 correlation with Vendian System, Australia
 18, 115–35
 event stratigraphy 126
 generalized time–space diagram, correlations
 between tectonic units 120
 naming 137–41
 etymology 140
 Waggoner's associations 139, 427
- Ediacaran–Cambrian boundary *see* Precambrian–
 Cambrian boundary
- Ediacaran–Ordovician, stratigraphic
 correlation 435–6
- Ediacaria*, morphology 300
- Ediacaria booleyi*
 five interpretations 291–3
 removal from genus *Ediacaria* 277–95
- Ediacaria flindersi* 279, 291
- Elatina Formation
 correlation with Smalfjord Formation,
 N Norway 139
 dating 140
 palaeomagnetic record 425–8
 rock magnetic experiments 71
- Elatina–Nuccaleena deglaciation 53–76
 Elatina Creek 61–2
 transition, key facies 58
 Trezona Bore 63, 65–7
- Enorama Creek, Ediacaran GSSP 54, 55–61,
 115–16
- Eoholynia* 272, 274
- Eotintinnopsis* (ciliated protozoan)
 Precambrian, China 151–6
 systematic palaeontology 155–6
- Ernietta*
 'colonies' 208, 213
 Namibia, Nama Group 197, 200
 quilting patterns 387–8, 400
 reconstruction 390
- Europe, Eastern, Vendian climatic indicators 16
- Fermeuse-style assemblages 299–307
- Finnmark
 correlations, Flinders Ranges 138
 Vendian climatic indicators 16
- Flinders Ranges 138, 425–8
 correlations with Finnmark, N Norway 138
 Elatina–Nuccaleena deglaciation 53–76
 location map 196
 Pound Subgroup
 Bonney Sandstone 122, 140, 197
 Rawnsley Quartzite 195–9, 426
- Flinders-style assemblages 299
- frondose organisms *see* *Charnia*; *Pambikalbae*
hasenohrae *gen. et sp.*
- Gangolihat Dolomite Formation, Lesser Himalaya,
 chemostratigraphy 83
- Garhwal, Lesser Himalaya,
 chemostratigraphy 79, 82, 84
- gastropod evolution 415–21
 main groups 415–18
 phylogenetic relationships 419–20
 shell forms, radiation 416
- Gawler Craton 425
- genes, regulatory genes in bilateral
 metazoans 372–4
- Georgina Basin
 carbon isotope trends 129
 location map 118
- Germany, Saxo-Thuringian Zone, Cadomian
 basement 35–51
- Glaessner, Martin 443–4
- Glaessnerina* 211, 212
 affinities 251
 gliding symmetry 399
- Glockerichnus* 5, 7
- Gondwana, Cadomian–Avalonian active
 margin 46
- Gymnoblastina* 217
- halite, sequestration as 'saline giants' 447
- Hallett Cove, Adelaide 68
- Helcionellidae 415–17
- Helminthoidichnites tenuis* 5, 7
- Helminthoraphe* 5, 7
- Hetang Formation, radiolarian
 skeletons 143–9
 stratigraphic column 144
 Yangtze Platform 423–4
- Hiemalora* 299, 300, 331–7
 reconstruction 335–6
 research history 331
 stratigraphic section 332
 taxonomy 336
- Himalaya, NW and NE Lesser
 Himalaya 77–101, 318–21
 Garhwal chemostratigraphy 79, 82, 84
 Krol Formation 77–82, 85–91, 318–21
 lithostratigraphy and palaeobiology 80
 Menga (Buxa) Limestone 93–5, 97–8, 81
 Mussoorie Syncline, isotope data 85–92
 Neoproterozoic Transition 88
 palaeoclimatic and biotic changes 88
 Ranjit Window 81

- Hormosiroidea* 226
honeycomb pattern, interpretations 343–9
conceptual paradigm 344–5
physical paradigm 345–7
inappropriate paradigms 347–9
Hox genes, regulatory genes in bilateral metazoans 372–4
Hydroactinidae 217
- Igarkiellidae 417
Indian subcontinent 77–101, 315–30, 439–41
Birmanian basin, Rajasthan 439–41
carbon isotope stratigraphy 439–41
Higher/Tethys Himalaya: Kashmir 321–2
Indo-Gangetic Plain 322–3
late Proterozoic biota 318–20
correlation charts 317, 325–6
geological setting 315–18
location map 314
Lesser Himalaya 77–101, 315–30, 318–21
Blaini–Krol–Tal Basin 79
Krol Belt 318–21
Mesoproterozoic–Neoproterozoic successions 325–6
NW and NE Lesser Himalaya 77–101
lithostratigraphy 80
palaeoclimatic and biotic changes 88
Peninsular India 323–8
Vendian climatic indicators 17
infaunal trace makers 406–9
continuous horizontal galleries in sand 406–7
correlation with small shelly fossils 410–12
cylindrical branching burrows 407–8
mud eaters 408–9
transitions in infaunal behaviour 409–10
Inkrylovia 208, 212
Iran (Takab area), Ediacaran–Ordovician microstratigraphy 433–7
correlation 435–6
Lalun Formation 434–5
petrography and stratigraphy 433–5
Ireland, Booley Bay Formation 277–95
age 288–91
location map 278
sedimentological setting 277–85
isotope data, Mussoorie Syncline, NW and NE Lesser Himalaya 89, 91–2
- Jakutsk, Lena River 408, 409
- Kessyusa Formation, Siberia 406–14
Khairkhanidae 418
Khorbusuonka River region, Siberia 406–14
Kimberella 157–79, 323, 369
future research directions 181–5
grazing tracks 174–7
morphology 160–74, 184
reconstructions 176, 185
stratigraphic setting 158
taphonomy and ecology 158–60, 182
King Island/NE Tasmania 129
generalized time–space diagram, correlations between tectonic units 120
Kotlin–Redkino, Vendian climatic zonation 20
Krol Formation, NW Himalaya 77–82, 85–91, 318–21
Baliana–Krol Group biota 319
Blaini–Krol–Tal Basin, carbon isotope chemostratigraphy 85–91
late Proterozoic successions 318–21
Mussoorie Syncline
isotope data 89
REE, Y, Sc and Th distribution 91
trace element distribution 90–1, 92
Kullingia (swing mark) 301
- Ladatheca annae* 411
Lakhanda Microbiota 431
Lalun Formation, Iran 434–5
Laplendian, Vendian climatic zonation 20–3
Lausitz Group 41–5
Leiosphaeridia jacutica/*L. crassa* Assemblage Zone 117, 119, 131, 132
Lena River, Jakutsk 408, 409
Louisella pedunculata priapulid 407
- Mackenzie, Vendian climatic indicators 16
magnetic susceptibility, temperature dependence 70–2
magnetics
cap dolostone 69–72
hysteresis loops 72
Verwey Transition 66
Manykodes (*Phycodes*) *pedum* 406, 411
Manykodes rectangulare 407–8
Marinoan glaciation
Australia 53, 126
cap carbonates, palaeomagnetism 54
Namibia 104
Mattaia 406–12
Mattaia miittensis burrows 406–7, 410
Mattaia tirasensis burrows 410–11
Mawsonites 299, 443
Medusinites 300
Menga Dolomite *see* Buxa Dolomite
metazoans
Bayesian analysis 357–9
before Cambrian explosion 369–71
bilateral 369–75
first appearance 126, 157, 447
regulatory genes 372–4
small-bodied crown phyla 371–2
early radiation 187
first appearance 447–8
origins 178
and origins of Animalia 447–8
paraphyly, eumetazoans and poriferans 356
phylogenetic relations 370
small-bodied crown phyla 371–2
trace fossils 371
microbial mats
demise 448
Lausitz group, Germany 41, 42
see also Vendobionts
microfossils
Buxa Dolomite, Lesser Himalaya 97–8
carbonaceous, White Sea Region 269–75
discoidal fossils 297–313
Hetang Formation, China 145–8

- organic-walled, Spain 230–2
- siliceous, China, earliest
 - Cambrian 146, 423–4
 - White Sea Region 269–75, 271
 - see also* small shelly fossils (SSF)
- Mila Formation, Iran 434, 435
- Mistaken Point, Newfoundland 301–6, 308
- Monomorphichnus lineatus* 5, 7
- Monomorphichnus lineatus*–*Treptichnus pedum* assemblage 232–3
- monoplacophorans, living 183
- Montes de Toledo–Guadalupe, Spain, Precambrian–Cambrian boundary 223–35
- Morphospace Analysis 377–86
 - formulation of morphospace for Ediacara biota 379–82
 - results 382–6
- Mussoorie Syncline
 - Blaini–Krol–Tal Basin, Lesser Himalaya
 - carbon isotope chemostratigraphy 85–91
 - trace elements 90, 91
 - isotope data, Krol Formation 89, 91–2
- Namalia*, quilting patterns 387–8
- Namibia, Nama Group
 - Ernietta* 197, 200, 208, 213, 387–8, 390, 400
 - preservation (taphonomic) processes 200
 - Preridinium* 402–4
- Namibia, Otavi Group 103–13
 - chemical profiles 108–10
 - Chuos Formation 104
 - location map 105
 - Rasthof Formation cap carbonates 27–34, 104–110
 - thin section data 107–8
- Namibian association 427
- Nantuo Formation, China 151–6
- Nasepia*, quilting patterns 387–8
- Nemakit–Daldynian
 - assemblages, Vendian climatic indicators 17, 18–20
 - carbon isotope chemostratigraphy 440
 - gastropod evolution 415–21
- Nemiana* 9, 158, 300
 - Russia and Ukraine, comparative taphonomy 263–5
- Neoneretes biserialis* 5, 7
- Neoneretes uniserialis* 5, 7
- Neoproterozoic glacials 447–8
 - ‘Snowball Earth’ hypothesis 21–3
- Neoproterozoic–Early Cambrian transition, Argentina (NW) 1–14
- Nereites saltensis* 5, 7
- Newfoundland
 - Avalon peninsula, *Charnia* fronds 237–57
 - Avalonian assemblages 303–7
 - Mistaken Point 301–6, 308
 - Vendian climatic indicators 16
- Ngalia Basin, location map 118
- Nimbia*
 - Booley Bay Formation 278, 285, 299
 - morphology 300
- Norway
 - Finnmark, Vendian climatic indicators 16
 - Smalfjord Formation, correlation with Elatina Formation, Australia 139
 - Stappogjedde Formation, Varanger, discoidal fossils 304–5
 - Vestertana Group, and Bunyeroo–Wonoka Formation transition 138
- Nuccaleena Formation cap dolostone 54, 59–61, 67–72, 120–1
 - boundary with Brachina Formation 121
 - dating 140
 - magnetics 69–72
- Officer Basin
 - acritarch zonation 119
 - carbon isotope trends 129
 - correlation with Adelaide Rift Complex 127
 - generalized time–space diagram, correlations between tectonic units 120
 - location map 118
 - sequence stratigraphy and seismic interpretation 126
 - succession 122–4
 - Oldhamia curvata* 5, 7
 - Oldhamia flabellata* 5, 7
 - Oldhamia radiata* 5, 7
 - Olenek River region, Siberia 406–14
 - Onuphionella*, mica-lined shelter 410
 - Onychochilidae 418
 - Oraparinna diapir 68
 - organic-walled microfossils 230–2
 - Oscillatoropsis* 269–70
 - Otavi Group *see* Namibia
 - Ovatoscutum* 217
 - oxygenation
 - and carbon isotope chemostratigraphy 362–3
 - Precambrian era 362–3
 - Shuram excursion 362
 - Precambrian–Cambrian boundary 431
 - Palaeolyngbia* 269–70
 - palaeopascichnids 262, 425
 - Palaeophycus* 5, 7, 8
 - Pambikalbae hasenohrae* *gen. et sp.* 195–222
 - biological relationships 207–18
 - living relatives 214–18
 - morphology 205–7
 - preservation 200–4
 - reconstruction 207, 215
 - systematic palaeontology 205
 - paraphyly
 - Cambrian ‘problematica’ 360–2
 - eumetazoans and poriferans 356
 - Precambrian oxygenation 362–3
 - pattern interpretations
 - englutinated fill skeletons 388
 - honeycomb 343–9
 - quilting 387–8
 - Pelagiellidae 418
 - Petalonamae 187–9, 209–18
 - ‘petalonamids’ *see* *Pambikalbae hasenohrae gen. et sp.*
 - phosphatic shells *see* small shelly fossils (SSF)
 - phosphogenesis, Precambrian–Cambrian boundary 439
 - Phyllozoon*
 - carpet-like recliners 389
 - quilting patterns 387, 400

- Phyllozoon hanseni* 212, 217
 reconstruction 209
 phylogenetics, Bayesian 356
 physonect siphonophores 215
Pilitella 272, 274
 placozoans 355
 plankton communities, early development and
 composition 148
 pneu architecture 387–8
Podolimirus 189
Podolodes burrows 406–11
 Podolodes tripleurum 409, 410
 Podolodes triplex 409
 Poland, Holy Cross mountains 407
 Porifera
 China, earliest Cambrian 424
 paraphyly 355–67
 Bayesian analysis 357–9
 Cambrian ‘problematica’ 360–2
 character acquisition 359–63
 spicules, preservation of radiolarian skeletons 146–8
 Precambrian era, oxygenation 362–3
 Precambrian–Cambrian boundary
 Flinders Ranges 426
 marine phosphogenesis 439
 origin of armour and infaunal shelters 405–14
 radiolarian skeletons 143–9
 Spain 223–35
 preservation (taphonomic) processes
 159, 182, 199–204
 priapulids 407–8
Primoflagella 430–1
Protolyellia 339–53
 affinities 351–2
 interpretation 342–51
 material 339–42
 reconstruction 349
 taphonomy 342, 349–51
 protozoans, Precambrian, China 151–6
Psammichnites 227–8, 406, 411–12
 non-*Psammichnites* 5, 7
 Psammocorallia 293, 339
 see also *Protolyellia*
Pteridinium carolinaense 198, 200, 208–9, 217, 320
 quilting patterns 387–8, 400
 specimens from Aar Farm, Namibia 402–4
 reconstructions 209, 391
 regeneration 393
 vanes 389
Pteridinium simplex 214
 reconstructions 391
 Puncovicana Formation, Argentina 1–14
 quilt structures, ‘Vendobionta’ 387–8, 399–404
 radiolarian skeletons, Early Cambrian 143–9
 coexistence with sponge spicules 147
 morphology 147–8
Radulichnus 181
 Rajasthan, Birmania basin 439–41
Rangea 210, 216, 390–3
 quilting patterns 388
 reconstructions 392
 vanes 389
 rangeomorphs 359–60
 Ranjit Window, Lesser Himalaya 81, 94
 Rasthof cap carbonates, Namibia 27–34, 103–13
 Rawnsley Quartzite (Pound Subgroup), Flinders
 Ranges 195–200, 426
 Breakfast Time Creek Member 197–8
 Chase Quartzite Member 197
 Ediacara Member 198–200
 Winnowie Member 198
 regulatory genes, bilateral metazoans 372–4
 Rheic Ocean 36
 ‘rhizocorallium’ spreites 408
 rhythmite sequences, cap carbonates 27–34
 Riphean animals 429–31
 Rodinia supercontinent 78
 break-up 78, 82–5
 modern continental outlines 139
Rokopelia *spp.* 183
 Rothstein Formation, backarc spreading 44–5, 47–8
Rugosoopsis 429–31
 Russia
 Ventogyrus 187–94, 208–14
 see also White Sea region
 sabelliditids 231, 232, 429
 ‘saline giants’ 447
 Saxo-Thuringian Zone (Cadmian basement)
 backarc basin 41–4
 geological map 36–8
 lithosections and geochronological data 38
 retroarc basin 45, 47–9
Sekwia 9, 300, 327
Selkirkia 4, 8
Serebrina 272, 274
 SHRIMP (sensitive high resolution ion microprobe)
 dating 125, 140
 Shuram excursion, oxygenation in Precambrian
 era 362
 Siberia
 Hiemalora 331–7
 Kessyusa Formation 406–14
 Khorbusuonka River region 406–14
 Olenek River region 406–14
 siliceous microfossils, China, earliest Cambrian
 146, 423–4
 Sinian association 437
 siphonophores 215–16, 218
 pneumatophore 216
Siphonophycus 269–70
Skolithos 158
 small shelly fossils (SSF)
 China, Precambrian–Cambrian boundary 143–9
 correlation with infaunal trace makers 410–12
 India 324
 Spain 230
 see also microfossils
 ‘Snowball Earth’ hypothesis 21–3
Sokolovina 429
 Soltanieh Formation, Iran 433–7
 South Africa, Vendian climatic indicators 17
 South American Andean Basin 3–14
 Spain, Precambrian–Cambrian boundary 223–35
 carbonaceous compressions 228–30
 geological setting 223–6

- location map 224
- organic-walled microfossils 230–2
- skeletal fossils 230
- stratigraphic nomenclature 223–6
- trace fossils 226–8
- sphaeromorphid acritarchs 324
- sponges *see* Porifera
- Sprigg, Reg C 443
- Spriggina* 300, 400
- Sr isotope data, Mussoorie Syncline, NW and NE
 - Lesser Himalaya 89, 91–2
- SSU rRNA molecule, phylogenetic relations 370
- Stannophyllum*, englutinated fill skeletons 388
- Stenotheceidae 417
- stromatolite assemblages
 - Australian Ediacaran System 129–30
 - Lesser Himalaya
 - Blaini–Krol–Tal sequence 85–7
 - Deoban–Gangolihat dolomite 78
 - Menga (Buxa) Limestone 93–5
 - Vindhyan Supergroup, India 324
- Stuart Shelf 120–2
 - acritarch zonation 119
- Sturtian glaciation, Namibia 104
- Sturtian (Umberatana) Group, Adelaide Rift Complex 116, 138
- Svalbard, Vendian climatic indicators 16
- Swarptuntia* 189, 200, 208, 212, 213, 217
 - upright form 389
- Sweden, Mickwitzia sandstone, *Protolyellia* 339–53
- Tanarium conoideum*/*Schizofusa risoria*/
Variomargosphaeridium litoschum
Assemblage Zone 119, 131, 132
- Tanarium irregulare*/*Ceratospaeridium glaberosum*/
Multifronsphaeridium pelorum Assemblage Zone
119, 131, 132
- taphonomic process 159, 182, 199–204
 - comparative taphonomy 259–67
 - Beltanelliformis* 260–2
 - Nemiana* 264
 - discoidal fossils 298–9
 - Ediacaria booleyi* 279
 - Protolyellia* 342, 349–51
- Tasmanadia cachii* 5, 7
- Tasmania, King Island 120, 129
- Tawuia* 435
- Tethys Himalayan sequence 77–8
- Thalassinoides* 5, 9
- Thaumaptilon* 212
- Tirasiana* 259–60, 299
 - morphology 300–1
- trace elements
 - Blaini–Krol–Tal Basin 90
 - Mussoorie Syncline 90–1, 92
- trace fossils 226–7, 371
 - infaunal trace makers 406–9
 - ‘Vendobionta’ 394–5
- Trenellidae 417
- Treptichnus* assemblage 224, 226–7
- Treptichnus cf. aequalternus* 5, 7, 8
- Trezona Bore
 - Brachina Formation 65–7, 69–70
 - Latina–Nuccaleena deglaciation 63, 65–7
- Trilobozoa 192
- Tuberlichnus* 226
- U–Pb aging, Newfoundland 427
- Ukraine
 - Beltanelliformis* 259–67
 - biostratigraphic correlation with White Sea Region 273
- Podolia
 - burrowing infauna 409–10
 - macrophyte algae 271–4
- Umberatana Group, Adelaide Rift Complex 116, 138
- Urals
 - Vendian climatic indicators 16
 - Vendian cyclic structure 22
- Uratanna Formation 426
- Vaizitsinia* 212
- Valdania* 212
- Veleropilina* 183
- Velumbrella* 282
- Vendian System and Period 427
 - climate zonation 20
 - climatic indicators 15–26
 - correlation with Ediacaran System 18
 - correlation of glacial horizons 16–17
 - macrofossil taxa 157
 - Russia
 - Beltanelliformis* 259–67
 - Nemiana* 158, 259–67
 - new data on *Kimberella* 157–79
 - Ventogyrus* 187–94
 - zonality 19
- ‘Vendobionta’ 211, 387–97
 - englutinated fill skeletons 388
 - quilt structures 387–8, 399–404
 - regeneration 393–4
 - resting traces 394–5
 - versatility 388–93
- Vendoconularia* 189
- Vendoglossa* 395, 396
- Vendotaenia*, Iran 435, 436
- Ventogyrus* 187–94, 208–9, 214
 - morphology 188–92, 208–9, 212
 - reconstructions 192–3
 - taxonomy 187–9
- ‘Verdun syndrome’ 405–14
- Vestertana Group, N Norway 138
 - and Bunyeroo–Wonoka Formation transition 138
- Wade, Mary 444
- Wearing Dolomite, Bunyeroo–Wonoka Formation transition 117, 121–2, 137–9
- White Sea Region
 - algal microfossils 271
 - Beltanelliformis* 259–67
 - biostratigraphic correlation with Podolia 273
 - carbonaceous microfossils 269–75
 - new data on *Kimberella* 157–79
 - Ventogyrus* 187–94
 - Verkhova Formation 198
 - Zimney Gory Formation 198, 262, 405
 - see also* Vendian System and Period
- Winnowie Member, Rawnsley Quartzite 198

- Wonoka Formation, Bunyeroo–Wonoka Formation
transition 121–2, 137–9, 426
- Wujiangella* (ciliated protozoan)
Precambrian, China 151–6
systematic palaeontology 155–6
- xenophyophores 387
- XRF analyser, scanning X-ray analytical microscope
(SXAM) 28–32
- Yakutsk, Lena River 408, 409
- Yangtze Plate, China
Hetang Shales 423–4
radiolarian skeletons 143–9
- Yarnemia* 214
- Yocheccionellidae 417
- Yonyangella* (ciliated protozoan)
Precambrian, China 151–6
systematic palaeontology 155–6
- Yorgia* 158, 448
quilting patterns 387–8, 400
resting traces 394
- Zaigun Formation, Iran 434
- Zimney Gory Formation, White Sea region 198, 262,
405–6
- zircons, SHRIMP dating 125, 140